

## Last time: change of coordinates in three-dimensions

### Theorem

Let  $T$  be a transformation from  $D \subset \mathbb{R}^3$  to  $\mathbb{R}^3$  such that

- $D$  and  $T(D)$  are “nice”;
- $\frac{\partial(x,y,z)}{\partial(u,v,w)} \neq 0$  and  $T$  is continuous on  $D$  except possibly on the boundary.

Suppose  $f$  is a continuous function on  $T(D)$ . Then

$$\iiint_{T(D)} f(x, y, z) dV_{xyz} = \iiint_D f(T(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| dV_{uvw}.$$

## Announcements

### **Participants Needed for a Research Study!**

*Message from PhD student Emily Teitelbaum*

My dissertation research seeks to contribute to higher education professionals' understanding of racism from the perspective of White American students.

The study will involve three 1-hour, in-person interviews with me. Participants who complete interviews will receive a \$10 gift card for each interview.

Are you:

- A full-time undergraduate student enrolled at UIUC;
- Self-identifying as White and American;
- At least 18 years old;
- Willing to be audio-recorded?

Please contact Emily Teitelbaum at [erteite2@illinois.edu](mailto:erteite2@illinois.edu) if interested!

## Practice with parametrizing surfaces

We just parametrized the sphere  $S = \{(x, y, z) \mid x^2 + y^2 + z^2 = 1\}$ .  
If we only want to parametrize the lower half of the sphere,  
 $S_h = \{(x, y, z) \mid x^2 + y^2 + z^2 = 1, z \leq 0\}$ , we should keep the  
same function  $\mathbf{r}(\phi, \theta)$ , but we should change the domain so that


- (a)  $\phi$  is in  $[0, \frac{\pi}{2}]$ ;
- (b)  $\phi$  is in  $[\frac{\pi}{2}, \pi]$ ;
- (c)  $\theta$  is in  $[0, \pi]$ ;
- (d)  $\theta$  is in  $[\pi, 2\pi]$ ;
- (e) I don't know.

Let  $D = \{(u, v) \mid 0 \leq u \leq 2\pi, -\frac{\pi}{2} \leq v \leq \frac{\pi}{2}\}$ .

Define  $\mathbf{r} : D \rightarrow \mathbb{R}^3$  by

$$\mathbf{r}(u, v) = (\sin v, \cos u \sin 2v, \sin u \sin 2v).$$

For any choice of  $v = v_0$ , the function  $u \mapsto \mathbf{r}(u, v_0)$  defines a curve on the surface  $S$ . What do these curves look like?



- (a) The curves which form circles on  $S$ .
- (b) The curves on  $S$  which pass through the origin  $(0, 0, 0)$ .
- (c) None of these.
- (d) Both of these.
- (e) I don't know.

## Practice with tangent planes

Find one or more partners. Parametrize the paraboloid  $z = x^2 + y^2$ .

Use the parametrization to find the tangent plane to the paraboloid at  $(1, 1, 2)$ .

- (a) We can't find a parametrization.
- (b) We found a parametrization  $\mathbf{r}(u, v)$ , but we can't find  $(u_0, v_0)$  such that  $\mathbf{r}(u_0, v_0) = (1, 1, 2)$ .
- (c) We found a parametrization and the point  $(u_0, v_0)$ , but we can't find the normal vector to the tangent plane.
- (d) We found the normal vector, but we can't remember how to write down the equation to the plane.
- (e) We did it!