

## Recall: integrating functions over curves

Let  $C$  be a smooth curve in  $\mathbb{R}^3$  parametrized by  $\mathbf{r}(t)$ ,  $a \leq t \leq b$ .  
Then

$$\text{Length}(C) = \int_a^b |\mathbf{r}'(t)| dt.$$

Furthermore, if  $g$  is a continuous function on  $C$ , then

$$\int_C g \, ds = \int_a^b g(\mathbf{r}(t)) |\mathbf{r}'(t)| dt.$$

**Today:** Integrate a function over a surface (or estimate the integral algebraically or geometrically).

**Applications:** Find surface area, mass, average value.

## Practice with surface area

Find the surface area of  $S = \{x^2 + y^2 + z^2 = 1\}$ .

### Step 1: Parametrize $S$

$$\mathbf{r}(\phi, \theta) = \langle \sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi \rangle,$$
$$0 \leq \phi \leq \pi; \quad 0 \leq \theta \leq 2\pi.$$

### Step 2: Calculate $|\mathbf{r}_\phi \times \mathbf{r}_\theta|$ .

$$\mathbf{r}_\phi(\phi, \theta) = \langle \cos \phi \cos \theta, \cos \phi \sin \theta, -\sin \phi \rangle;$$
$$\mathbf{r}_\theta(\phi, \theta) = \langle -\sin \phi \sin \theta, \sin \phi \cos \theta, 0 \rangle.$$

So

$$\begin{aligned}\mathbf{r}_\phi \times \mathbf{r}_\theta &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \cos \phi \cos \theta & \cos \phi \sin \theta & -\sin \phi \\ -\sin \phi \sin \theta & \sin \phi \cos \theta & 0 \end{vmatrix} \\ &= \mathbf{i}(\sin^2 \phi \cos \theta) - \mathbf{j}(-\sin^2 \phi \sin \theta) \\ &\quad + \mathbf{k}(\sin \phi \cos \phi \cos^2 \theta + \sin \phi \cos \phi \sin^2 \theta) \\ &= \langle \sin^2 \phi \cos \theta, \sin^2 \phi \sin \theta, \sin \phi \cos \phi \rangle.\end{aligned}$$

Therefore

$$\begin{aligned}|\mathbf{r}_\phi \times \mathbf{r}_\theta| &= \sqrt{\sin^4 \phi (\cos^2 \theta + \sin^2 \theta) + \sin^2 \phi \cos^2 \phi} \\ &= \sqrt{\sin^2 \phi (\sin^2 \phi + \cos^2 \phi)} \\ &= \sqrt{\sin^2 \phi} = \sin \phi\end{aligned}$$

(since  $\sin \phi \geq 0$  on  $D$ ).

So the surface area of the sphere is

$$\begin{aligned}\iint_D |\mathbf{r}_\phi \times \mathbf{r}_\theta| dA &= \int_0^{2\pi} \int_0^\pi \sin \phi d\phi d\theta \\ &= 2\pi [-\cos \phi]_0^\pi \\ &= 4\pi.\end{aligned}$$

## Practice with surface area

Consider a can with sides given by the cylinder  $\{x^2 + y^2 = 1, -1 \leq z \leq 1\}$ , parametrized by

$$\mathbf{r}(\theta, z) = \langle \cos \theta, \sin \theta, z \rangle,$$
$$0 \leq \theta \leq 2\pi, \quad -1 \leq z \leq 1.$$

Find the surface area of the can. (Don't forget the top and bottom!)

- (a)  $2\pi$
- (b)  $4\pi$
- (c)  $6\pi$
- (d)  $8\pi$
- (e) I don't know how.