

CATEGORIFICATION AND HEISENBERG DOUBLES ARISING FROM TOWERS OF ALGEBRAS

ALISTAIR SAVAGE AND ODED YACOBI

ABSTRACT. The Grothendieck groups of the categories of finitely generated modules and finitely generated projective modules over a tower of algebras can be endowed with (co)algebra structures that, in many cases of interest, give rise to a dual pair of Hopf algebras. Moreover, given a dual pair of Hopf algebras, one can construct an algebra called the Heisenberg double, which is a generalization of the classical Heisenberg algebra. The aim of this paper is to study Heisenberg doubles arising from towers of algebras in this manner. First, we develop the basic representation theory of such Heisenberg doubles and show that if induction and restriction satisfy Mackey-like isomorphisms then the Fock space representation of the Heisenberg double has a natural categorification. This unifies the existing categorifications of the polynomial representation of the Weyl algebra and the Fock space representation of the Heisenberg algebra. Second, we develop in detail the theory applied to the tower of 0-Hecke algebras, obtaining new Heisenberg-like algebras that we call *quasi-Heisenberg algebras*. As an application of a generalized Stone–von Neumann Theorem, we give a new proof of the fact that the ring of quasisymmetric functions is free over the ring of symmetric functions.

CONTENTS

1. Introduction	2
2. The Heisenberg double	4
3. Towers of algebras and categorification of Fock space	8
4. Hecke-like towers of algebras	16
5. Nilcoxeter algebras	20
6. Hecke algebras at generic parameters	21
7. Hecke algebras at roots of unity	23
8. 0-Hecke algebras	24
9. Application: QSym is free over Sym	28
References	29

Date: September 10, 2013.

2010 *Mathematics Subject Classification.* 16D90, 16G10, 16T05.

Key words and phrases. Heisenberg double, tower of algebras, categorification, Hopf algebra, Hecke algebra, quasisymmetric function, noncommutative symmetric function, Heisenberg algebra, Fock space.

The first author was supported by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada. The second author was supported by an AMS–Simons Travel Grant and the University of Toronto.

1. INTRODUCTION

The interplay between symmetric groups and the Heisenberg algebra has a rich history, with implications in combinatorics, representation theory, and mathematical physics. A foundational result in this theory is due to Geissinger, who gave a representation theoretic realization of the bialgebra of symmetric functions Sym by considering the Grothendieck groups of representations of all symmetric groups over a field \mathbb{k} of characteristic zero (see [Gei77]). In particular, he constructed an isomorphism of bialgebras

$$\text{Sym} \cong \bigoplus_{n=0}^{\infty} \mathcal{K}_0(\mathbb{k}[S_n]\text{-mod}),$$

where $\mathcal{K}_0(\mathcal{C})$ denotes the Grothendieck group of an abelian category \mathcal{C} . Multiplication is described by the induction functor

$$[\text{Ind}] : \mathcal{K}_0(\mathbb{k}[S_n]\text{-mod}) \otimes \mathcal{K}_0(\mathbb{k}[S_m]\text{-mod}) \rightarrow \mathcal{K}_0(\mathbb{k}[S_{n+m}]\text{-mod}),$$

while comultiplication is given by restriction. Mackey theory for induction and restriction in symmetric groups implies that the coproduct is an algebra homomorphism. For each S_n -module V , multiplication by the class $[V] \in \mathcal{K}_0(\mathbb{k}[S_n]\text{-mod})$ defines an endomorphism of $\bigoplus_{n=0}^{\infty} \mathcal{K}_0(\mathbb{k}[S_n]\text{-mod})$. These endomorphisms, together with their adjoints, define a representation of the Heisenberg algebra on $\bigoplus_{n=0}^{\infty} \mathcal{K}_0(\mathbb{k}[S_n]\text{-mod})$.

Geissinger's construction was q -deformed by Zelevinsky in [Zel81], who replaced the group algebra of the symmetric group $\mathbb{k}[S_n]$ by the Hecke algebra $H_n(q)$ at generic q . Again, endomorphisms of the Grothendieck group given by multiplication by classes $[V]$, together with their adjoints, generate a representation of the Heisenberg algebra.

The above results can be enhanced to a categorification of the Heisenberg algebra and its Fock space representation via categories of modules over symmetric groups and Hecke algebras. A strengthened version of this categorification, which includes information about the natural transformations involved, was given in [Kho] for the case of symmetric groups and in [LS13] for the case of Hecke algebras.

The group algebras of symmetric groups and Hecke algebras are both examples of *towers of algebras*. A tower of algebras is a graded algebra $A = \bigoplus_{n \geq 0} A_n$, where each A_n is itself an algebra (with a different multiplication) and such that the multiplication in A induces homomorphisms $A_m \otimes A_n \rightarrow A_{m+n}$ of algebras (see Definition 3.1). In addition to those mentioned above, examples include nilcoxeter algebras, 0-Hecke algebras, Hecke algebras at roots of unity, wreath products (semidirect products of symmetric groups and finite groups, see [FJW00, CL12]), group algebras of finite general linear groups, and cyclotomic Khovanov–Lauda–Rouquier algebras (quiver Hecke algebras). To a tower of algebras, one can associate the \mathbb{Z} -modules $\mathcal{G}(A) = \bigoplus_n \mathcal{K}_0(A_n\text{-mod})$ and $\mathcal{K}(A) = \bigoplus_n \mathcal{K}_0(A_n\text{-pmod})$, where $A_n\text{-mod}$ (respectively $A_n\text{-pmod}$) is the category of finitely generated (respectively finitely generated projective) A_n -modules. In many cases, induction and restriction endow $\mathcal{K}(A)$ and $\mathcal{G}(A)$ with the structure of dual Hopf algebras. For example, in [BL09] Bergeron and Li introduced a set of axioms for a tower of algebras that ensure this duality (although the axioms we consider in the current paper are different).

One of the main goals of the current paper is to generalize the above categorifications of the Fock space representation of the Heisenberg algebra to more general towers of algebras. We see that, in the general situation, the Heisenberg algebra \mathfrak{h} is replaced by the *Heisenberg double* (see Definition 2.6) of $\mathcal{G}(A)$. The Heisenberg double of a Hopf algebra is different

from, but closely related to, the more well known Drinfeld quantum double. As a \mathbb{k} -module, the Heisenberg double $\mathfrak{h}(H^+, H^-)$ of a Hopf algebra H^+ (over \mathbb{k}) with dual H^- is isomorphic to $H^+ \otimes_{\mathbb{k}} H^-$, and the factors H^- and H^+ are subalgebras. The most well known example of this construction is when H^- and H^+ are both the Hopf algebra of symmetric functions, which is self-dual. In this case the Heisenberg double is the classical Heisenberg algebra. In general, there is a natural action of $\mathfrak{h}(H^+, H^-)$ on its Fock space H^+ generalizing the usual Fock space representation of the Heisenberg algebra. Our first result (Theorem 2.11) is a generalization of the well known Stone–von Neumann Theorem to this Heisenberg double setting.

In the special case of dual Hopf algebras arising from a tower of algebras A , we denote the Heisenberg double by $\mathfrak{h}(A)$ and the resulting Fock space by $\mathcal{F}(A)$. In this situation, there is a natural subalgebra of $\mathfrak{h}(A)$. In particular, the image $\mathcal{G}_{\text{proj}}(A)$ of the natural Cartan map $\mathcal{K}(A) \rightarrow \mathcal{G}(A)$ is a Hopf subalgebra of $\mathcal{G}(A)$, and we consider also the *projective* Heisenberg double $\mathfrak{h}_{\text{proj}}(A)$ which is, by definition, the subalgebra of $\mathfrak{h}(A)$ generated by $\mathcal{G}_{\text{proj}}(A)$ and $\mathcal{K}(A)$. Then $\mathfrak{h}_{\text{proj}}(A)$ acts on its Fock space $\mathcal{G}_{\text{proj}}(A)$, and a Stone–von Neumann type theorem also holds for this action (see Proposition 3.15).

We then focus our attention on towers of algebras that satisfy natural compatibility conditions between induction and restriction analogous to the well known Mackey theory for finite groups. We call these towers of algebras *strong* (see Definition 3.4) and give a necessary and sufficient condition for them to give rise to dual pairs of Hopf algebras (i.e. be *dualizing*). Our central theorem (Theorem 3.18) is that, for such towers, the Fock spaces representations of the algebras $\mathfrak{h}(A)$ and $\mathfrak{h}_{\text{proj}}(A)$ admit categorifications coming from induction and restrictions functors on $\bigoplus_n A_n$ -mod and $\bigoplus_n A_n$ -pmod respectively.

To illustrate our main result, we apply it to several towers of algebras that are quotients of group algebras of braid groups by quadratic relations (see Definition 4.1). We first show that all towers of this form are strong and dualizing. Examples include the towers of nilcoxeter algebras, Hecke algebras, and 0-Hecke algebras. Starting with the tower of nilcoxeter algebras, we recover Khovanov’s categorification of the polynomial representation of the Weyl algebra (see Section 5). Taking instead the tower of Hecke algebras at a generic parameter, we recover (weakened versions of) the categorifications of the Fock space representation of the Heisenberg algebra described by Khovanov and Licata–Savage (see Section 6). Considering the tower of Hecke algebras at a root of unity, we obtain a different categorification of the Fock space representation of the Heisenberg algebra (Proposition 7.2) which, in the setting of the existing categorification of the basic representation of affine \mathfrak{sl}_n using this tower, corresponds to the principal Heisenberg subalgebra. In this way, we see that Theorem 3.18 provides a uniform treatment and generalization of these categorification results. A major feature of our categorification is that it does not depend on any presentation of the algebras in question, in contrast to many categorification results in the literature.

We explore the example of the tower A of 0-Hecke algebras in some detail. In this case, it is known that $\mathcal{K}(A)$ and $\mathcal{G}(A)$ are the Hopf algebras of noncommutative symmetric functions and quasisymmetric functions respectively. However, the algebras $\mathfrak{h}(A)$ and $\mathfrak{h}_{\text{proj}}(A)$, which we call the *quasi-Heisenberg algebra* and *projective quasi-Heisenberg algebra*, do not seem to have been studied in the literature. We give presentations of these algebras by generators and relations (see Section 8.4). The algebra $\mathfrak{h}_{\text{proj}}(A)$ turns out to be particularly simple as it

is a “de-abelianization” of the usual Heisenberg algebra (see Proposition 8.7). As an application of the generalized Stone–von Neumann Theorem in this case, we give a representation theoretic proof of the fact that the ring of quasisymmetric functions is free over the ring of symmetric functions (Proposition 9.2). Our proof is quite different than previous proofs appearing in the literature.

There are many more examples of towers of algebras for which we do not work out the detailed implications of our main theorem. Furthermore, we expect that the results of this paper could be generalized to apply to towers of superalgebras. Examples of such towers include Sergeev algebras and 0-Hecke-Clifford algebras. We leave such generalizations for future work.

Notation. We let \mathbb{N} and \mathbb{N}_+ denote the set of nonnegative and positive integers respectively. We let \mathbb{k} be a commutative ring (with unit) and \mathbb{F} be a field. For $n \in \mathbb{N}$, we let $\mathcal{P}(n)$ denote the set of all partitions of n , with the convention that $\mathcal{P}(0) = \{\emptyset\}$, and let $\mathcal{P} = \bigcup_{n \in \mathbb{N}} \mathcal{P}(n)$. Similarly, we let $\mathcal{C}(n)$ denote the set of all compositions of n and let $\mathcal{C} = \bigcup_{n \in \mathbb{N}} \mathcal{C}(n)$. For a composition or partition α , we let $\ell(\alpha)$ denote the length of α (i.e. the number of nonzero parts) and let $|\alpha|$ denote its size (i.e. the sum of its parts). By a slight abuse of terminology, we will use the terms *module* and *representation* interchangeably.

Acknowledgements. The authors would like to thank N. Bergeron, J. Bernstein, M. Khovanov, A. Lauda, A. Licata, J.-Y. Thibon, and M. Zabrocki for useful conversations.

2. THE HEISENBERG DOUBLE

In this section, we review the definition of the Heisenberg double of a Hopf algebra and state some important facts about its natural Fock space representation. In particular, we prove a generalization of the well known Stone–von Neumann Theorem (Theorem 2.11).

We fix a commutative ring \mathbb{k} and all algebras, coalgebras, bialgebras and Hopf algebras will be over \mathbb{k} . We will denote the multiplication, comultiplication, unit, counit and antipode of a Hopf algebra by ∇ , Δ , η , ε and S respectively. We will use Sweedler notation

$$\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}$$

for coproducts. For a \mathbb{k} -module V , we will simply write $\text{End } V$ for $\text{End}_{\mathbb{k}} V$. All tensor products are over \mathbb{k} unless otherwise indicated.

2.1. Dual Hopf algebras. We begin by recalling the notion of dual (graded connected) Hopf algebras.

Definition 2.1 (Graded connected Hopf algebra). We say that a bialgebra H is *graded* if $H = \bigoplus_{n \in \mathbb{N}} H_n$, where each H_n , $n \in \mathbb{N}$, is finitely generated and free as a \mathbb{k} -module, and the following conditions are satisfied:

$$\begin{aligned} \nabla(H_k \otimes H_\ell) &\subseteq H_{k+\ell}, & \Delta(H_k) &\subseteq \bigoplus_{j=0}^k H_j \otimes H_{k-j}, & k, \ell \in \mathbb{N}, \\ \eta(\mathbb{k}) &\subseteq H_0, & \varepsilon(H_k) &= 0 \text{ for } k \in \mathbb{N}_+. \end{aligned}$$

We say that H is *graded connected* if it is graded and $H_0 = \mathbb{k}1_H$. Recall that a graded connected bialgebra is a Hopf algebra with invertible antipode (see, for example, [Haz08, p. 389, Cor. 5]) and thus we will also call such an object a *graded connected Hopf algebra*.

If $H = \bigoplus_{n \in \mathbb{N}} H_n$ is a graded bialgebra, then its *graded dual* $\bigoplus_{n \in \mathbb{N}} H_n^*$ is also a graded bialgebra.

Remark 2.2. In general, one need not assume that the H_n are free as \mathbb{k} -modules. Instead, one needs only assume that

$$(2.1) \quad H_k^* \otimes H_\ell^* \cong (H_k \otimes H_\ell)^* \text{ for all } k, \ell \in \mathbb{N}$$

in order for the graded dual to be a graded bialgebra. However, since our interest lies mainly in dual Hopf algebras arising from towers of algebras, for which the H_n are free as \mathbb{k} -modules, we will make this assumption from the start (in which case (2.1) is automatically satisfied).

Definition 2.3 (Hopf pairing). If H and H' are Hopf algebras over \mathbb{k} , then a *Hopf pairing* is a bilinear map $\langle \cdot, \cdot \rangle: H \times H' \rightarrow \mathbb{k}$ such that

$$\begin{aligned} \langle ab, x \rangle &= \langle a \otimes b, \Delta(x) \rangle = \sum_{(x)} \langle a, x_{(1)} \rangle \langle b, x_{(2)} \rangle, \\ \langle a, xy \rangle &= \langle \Delta(a), x \otimes y \rangle = \sum_{(a)} \langle a_{(1)}, x \rangle \langle a_{(2)}, y \rangle, \\ \langle 1_H, x \rangle &= \varepsilon(x), \quad \langle a, 1_{H'} \rangle = \varepsilon(a), \end{aligned}$$

for all $a, b \in H$, $x, y \in H'$. Note that such a Hopf pairing automatically satisfies $\langle a, S(x) \rangle = \langle S(a), x \rangle$ for all $a \in H$ and $x \in H'$.

Recall that, for \mathbb{k} -modules V and W , a bilinear form $\langle \cdot, \cdot \rangle: V \times W \rightarrow \mathbb{k}$ is called a *perfect pairing* if the induced map $\Phi: V \rightarrow W^*$ given by $\Phi(v)(w) = \langle v, w \rangle$ is an isomorphism.

Definition 2.4 (Dual pair). We say that (H^+, H^-) is a *dual pair* of Hopf algebras if H^+ and H^- are both graded connected Hopf algebras and H^\pm is graded dual to H^\mp (as a Hopf algebra) via a perfect Hopf pairing $\langle \cdot, \cdot \rangle: H^- \times H^+ \rightarrow \mathbb{k}$.

2.2. The Heisenberg double. For the remainder of this section, we fix a dual pair (H^+, H^-) of Hopf algebras. Then any $a \in H^+$ defines an element ${}^L a \in \text{End } H^+$ by left multiplication. Similarly, any $x \in H^-$ defines an element ${}^R x \in \text{End } H^-$ by right multiplication, whose adjoint ${}^R x^*$ is an element of $\text{End } H^+$. (In the case that H^+ or H^- is commutative, we often omit the superscript L or R .) In this way we have \mathbb{k} -algebra homomorphisms

$$(2.2) \quad H^+ \hookrightarrow \text{End } H^+, \quad a \mapsto {}^L a,$$

$$(2.3) \quad H^- \hookrightarrow \text{End } H^+, \quad x \mapsto {}^R x^*.$$

The action of H^- on H^+ given by (2.3) is called the *left regular action*.

Lemma 2.5. *The left regular action of H^- on H^+ is given by*

$${}^R x^*(a) = \sum_{(a)} \langle x, a_{(2)} \rangle a_{(1)} \quad \text{for all } x \in H^-, a \in H^+.$$

Proof. For all $x, y \in H^-$ and $a \in H^+$, we have

$$\begin{aligned} \langle y, {}^R x^*(a) \rangle &= \langle yx, a \rangle = \langle y \otimes x, \Delta(a) \rangle \\ &= \sum_{(a)} \langle y \otimes x, a_{(1)} \otimes a_{(2)} \rangle = \sum_{(a)} \langle y, a_{(1)} \rangle \langle x, a_{(2)} \rangle = \langle y, \sum_{(a)} \langle x, a_{(2)} \rangle a_{(1)} \rangle. \end{aligned}$$

The result then follows from the nondegeneracy of the bilinear form. \square

It is clear that the map (2.2) is injective. The map (2.3) is also injective. Indeed, for $x \in H^-$, choose $a \in H^+$ such that $\langle x, a \rangle \neq 0$. Then $\langle 1, {}^R x^*(a) \rangle = \langle x, a \rangle \neq 0$, and so ${}^R x^* \neq 0$.

Since $H^+ = \bigoplus_{n \in \mathbb{N}} H_n^+$ is \mathbb{N} -graded, we have a natural algebra \mathbb{Z} -grading $\text{End } H^+ = \bigoplus_{n \in \mathbb{Z}} \text{End}_n H^+$. It is routine to verify that the map (2.2) sends H_n^+ to $\text{End}_n H^+$ and the map (2.3) sends H_n^- to $\text{End}_{-n} H^+$ for all $n \in \mathbb{N}$.

Definition 2.6 (The Heisenberg double, [STS94, Def. 3.1]). We define $\mathfrak{h}(H^+, H^-)$ to be the *Heisenberg double* of H^+ . More precisely $\mathfrak{h}(H^+, H^-) \cong H^+ \otimes H^-$ as \mathbb{k} -modules, and we write $a \# x$ for $a \otimes x$, $a \in H^+$, $x \in H^-$, viewed as an element of $\mathfrak{h}(H^+, H^-)$. Multiplication is given by

$$(2.4) \quad (a \# x)(b \# y) := \sum_{(x)} a {}^R x_{(1)}^*(b) \# x_{(2)} y = \sum_{(x), (b)} \langle x_{(1)}, b_{(2)} \rangle a b_{(1)} \# x_{(2)} y.$$

We will often view H^+ and H^- as subalgebras of $\mathfrak{h}(H^+, H^-)$ via the maps $a \mapsto a \# 1$ and $x \mapsto 1 \# x$ for $a \in H^+$ and $x \in H^-$. Then we have $ax = a \# x$. When the context is clear, we will simply write \mathfrak{h} for $\mathfrak{h}(H^+, H^-)$. We have a natural grading $\mathfrak{h} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{h}_n$, where $\mathfrak{h}_n = \bigoplus_{k-\ell=n} H_k^+ \# H_\ell^-$.

Remark 2.7. The Heisenberg double is a twist of the Drinfeld quantum double by a right 2-cocycle (see [Lu94, Th. 6.2]).

Lemma 2.8. *If $x \in H^-$ and $a, b \in H^+$, then*

$${}^R x^*(ab) = \sum_{(x)} {}^R x_{(1)}^*(a) {}^R x_{(2)}^*(b).$$

Proof. For $x, y \in H^-$ and $a, b \in H^+$, we have

$$\begin{aligned} \langle y, {}^R x^*(ab) \rangle &= \langle yx, ab \rangle = \langle \Delta(yx), a \otimes b \rangle = \langle \Delta(y)\Delta(x), a \otimes b \rangle \\ &= \langle \Delta(y), {}^R \Delta(x)^*(a \otimes b) \rangle = \langle \Delta(y), \sum_{(x)} {}^R x_{(1)}^*(a) \otimes {}^R x_{(2)}^*(b) \rangle = \langle y, \sum_{(x)} {}^R x_{(1)}^*(a) {}^R x_{(2)}^*(b) \rangle. \end{aligned}$$

The result then follows from the nondegeneracy of the bilinear form. \square

Interchanging H^- and H^+ in the construction of the Heisenberg double results in the opposite algebra: $\mathfrak{h}(H^-, H^+) \cong \mathfrak{h}(H^+, H^-)^{\text{op}}$ (see [Lu94, Prop. 5.3]).

2.3. Fock space. We now introduce a natural representation of the algebra \mathfrak{h} .

Definition 2.9 (Vacuum vector). An element v of an \mathfrak{h} -module V is called a *lowest weight* (resp. *highest weight*) *vacuum vector* if $\mathbb{k}v \cong \mathbb{k}$ and $H^-v = 0$ (resp. $H^+v = 0$).

Definition 2.10 (Fock space). The algebra \mathfrak{h} has a natural (left) representation on H^+ given by

$$(a \# x)(b) = a {}^R x^*(b), \quad a, b \in H^+, \quad x \in H^-.$$

We call this the *lowest weight Fock space representation* of $\mathfrak{h}(H^+, H^-)$ and denote it by $\mathcal{F} = \mathcal{F}(H^+, H^-)$. Note that this representation is generated by the lowest weight vacuum vector $1 \in H^+$.

Suppose X^+ is a graded subalgebra of H^+ that is invariant under the left regular action of H^- on H^+ . Then $X^+ \# H^-$ is a subalgebra of $\mathfrak{h}(H^+, H^-)$ acting naturally on X^+ . The following result (when $X^+ = H^+$) is a generalization of the Stone–von Neumann Theorem to the setting of an arbitrary Heisenberg double.

Theorem 2.11. *Let X^+ be a subalgebra of H^+ that is invariant under the left regular action of H^- on H^+ .*

- (a) *The only $(X^+\#H^-)$ -submodules of X^+ are those of the form IX^+ for some ideal I of \mathbb{k} . In particular, if \mathbb{k} is a field, then X^+ is irreducible as an $(X^+\#H^-)$ -module.*
- (b) *Let $\mathbb{k}^- \cong \mathbb{k}$ (isomorphism of \mathbb{k} -modules) be the representation of H^- such that H_n^- acts as zero for all $n > 0$ and $H_0^- \cong \mathbb{k}$ acts by left multiplication. Then X^+ is isomorphic to the induced module $\text{Ind}_{H^-}^{X^+\#H^-} \mathbb{k}^-$ as an $(X^+\#H^-)$ -module.*
- (c) *Any $(X^+\#H^-)$ -module generated by a lowest weight vacuum vector is isomorphic to X^+ .*

If $X^+ = H^+$ then $X^+\#H^- = \mathfrak{h}(H^+, H^-)$ and the module X^+ is the lowest weight Fock space \mathcal{F} . In that case we also have the following.

- (d) *The lowest weight Fock space representation \mathcal{F} of \mathfrak{h} is faithful.*

Proof. (a) Clearly, if I is an ideal of \mathbb{k} , then IX^+ is a submodule of X^+ . Now suppose $W \subseteq X^+$ is a nonzero submodule, and let

$$I = \{c \in \mathbb{k} \mid c1 \in W\}.$$

It is easy to see that I is an ideal of \mathbb{k} . We claim that $W = IX^+$. Since the element 1 generates X^+ , we clearly have $IX^+ \subseteq W$. Now suppose there exists $a \in W$ such that $a \notin IX^+$. Without loss of generality, we can write $a = \sum_{n=0}^{\ell} a_n$ for $a_n \in H_n^+$, $\ell \in \mathbb{N}$, and $a_\ell \notin IX^+$ (otherwise, consider $a - a_\ell$). Let b_1, \dots, b_m be a basis of H_ℓ^+ such that b_1, \dots, b_k is a basis of X_ℓ^+ , for $k = \dim_{\mathbb{k}} X_\ell^+$. Let x_1, \dots, x_m be the dual basis of H_ℓ^- . Then it is easy to verify that $\sum_{j=1}^k b_j \# x_j$ acts as the identity on X_ℓ^+ and as zero on X_n^+ for $n < \ell$. Thus,

$$a_\ell = \sum_{j=1}^k (b_j \# x_j)(a) \in \sum_{j=1}^k b_j IX^+ \subseteq IX^+,$$

since ${}^R x_j^*(a) \in W \cap H_0^+$ for all $j = 1, \dots, k$. This contradiction completes the proof.

- (b) We have an injective homomorphism of H^- -modules

$$\mathbb{k}^- \hookrightarrow \text{Res}_{H^-}^{X^+\#H^-} X^+, \quad 1 \mapsto 1.$$

Since induction is left adjoint to restriction (see, for example, [CR81, (2.19)]), this gives rise to a homomorphism of \mathfrak{h} -modules

$$(2.5) \quad \text{Ind}_{H^-}^{X^+\#H^-} \mathbb{k}^- \rightarrow X^+, \quad 1 \mapsto 1.$$

Since the element 1 generates X^+ , this map is surjective. Now,

$$\text{Ind}_{H^-}^{X^+\#H^-} \mathbb{k}^- = (X^+\#H^-) \otimes_{H^-} \mathbb{k}^- = X^+ H^- \otimes_{H^-} \mathbb{k}^-.$$

It follows that, as a left X^+ -module, $\text{Ind}_{H^-}^{X^+\#H^-} \mathbb{k}^-$ is a quotient of $X^+ \otimes_{\mathbb{k}} \mathbb{k}^- \cong X^+$ and the map (2.5) is the identity map, hence an isomorphism.

(c) Suppose V is a representation of $X^+\#H^-$ generated by a lowest weight vacuum vector v_0 . Then, as above, we have an injective homomorphism of H^- -modules

$$\mathbb{k}^- \hookrightarrow \text{Res}_{H^-}^{X^+\#H^-} V, \quad 1 \mapsto v_0,$$

and thus a homomorphism of $(X^+\#H^-)$ -modules

$$(2.6) \quad \text{Ind}_{H^-}^{X^+\#H^-} \mathbb{k}^- \rightarrow V, \quad 1 \mapsto v_0.$$

Since V is generated by v_0 , this map is surjective. Since $\text{Ind}_{H^-}^{X^+ \# H^-} \mathbb{k}^- \cong X^+$, it follows easily from part (a) that it is also injective.

(d) Suppose α is a nonzero element of \mathfrak{h} . Write $\alpha = \alpha' + \alpha''$ where α' is a nonzero element of $H^+ \# H_n^-$ for some $n \in \mathbb{N}$ and $\alpha'' \in \sum_{k>n} H^+ \# H_k^-$. Choose a basis x_1, \dots, x_m of H_n^- and let b_1, \dots, b_m denote the dual basis of H_n^+ . Then we can write $\alpha' = \sum_{j=1}^m a_j \# x_j$ for some $a_j \in H^+$. Since $\alpha' \neq 0$, we have $a_j \neq 0$ for some j . Then $\alpha(b_j) = \alpha'(b_j) = a_j \neq 0$. Thus the action of \mathfrak{h} on \mathcal{F} is faithful. \square

Remark 2.12. By Theorem 2.11(d), we may view \mathfrak{h} as the subalgebra of $\text{End } H^+$ generated by ${}^L a$, $a \in H^+$, and ${}^R x^*$, $x \in H^-$.

3. TOWERS OF ALGEBRAS AND CATEGORIFICATION OF FOCK SPACE

In this section, we consider dual Hopf algebras arising from towers of algebras. In this case, we are able to deduce some further results about the Heisenberg double \mathfrak{h} . We then prove our main result, that towers of algebras give rise to categorifications of the lowest weight Fock space representation of \mathfrak{h} (Theorem 3.18). Recall that \mathbb{F} is an arbitrary field.

3.1. Modules categories and their Grothendieck groups. For an arbitrary \mathbb{F} -algebra B , let $B\text{-mod}$ denote the category of finitely generated left B -modules and let $B\text{-pmod}$ denote the category of finitely generated projective left B -modules. We then define

$$G_0(B) = \mathcal{K}_0(B\text{-mod}) \quad \text{and} \quad K_0(B) = \mathcal{K}_0(B\text{-pmod}),$$

where $\mathcal{K}_0(\mathcal{C})$ denotes the Grothendieck group of an abelian category \mathcal{C} . We denote the class of an object $M \in \mathcal{C}$ in $\mathcal{K}_0(\mathcal{C})$ by $[M]$. Note that since all short exact sequences in $B\text{-pmod}$ split, $K_0(B)$ is also the split Grothendieck group of $B\text{-pmod}$.

There is a natural bilinear form

$$(3.1) \quad \langle \cdot, \cdot \rangle: K_0(B) \otimes G_0(B) \rightarrow \mathbb{Z}, \quad \langle [P], [M] \rangle = \dim_{\mathbb{F}} \text{Hom}_B(P, M).$$

If B is a finite dimensional algebra, let V_1, \dots, V_s be a complete list of nonisomorphic simple B -modules. If P_i is the projective cover of V_i for $i = 1, \dots, s$, then P_1, \dots, P_s is a complete list of nonisomorphic indecomposable projective B -modules (see, for example, [ARS95, Cor. I.4.5]) and we have

$$G_0(B) = \bigoplus_{i=1}^s \mathbb{Z}[V_i] \quad \text{and} \quad K_0(B) = \bigoplus_{i=1}^s \mathbb{Z}[P_i].$$

If \mathbb{F} is algebraically closed, then

$$(3.2) \quad \langle [P_i], [V_j] \rangle = \delta_{ij} \quad \text{for } 1 \leq i, j \leq s,$$

and so the pairing (3.1) is perfect.

Suppose $\varphi: B \rightarrow A$ is an algebra homomorphism. Then we can consider A as a left B -module via the action $(b, a) \mapsto \varphi(b)a$. Similarly, we can consider A as a right B -module. Then we have *induction* and *restriction* functors

$$\text{Ind}_B^A: B\text{-mod} \rightarrow A\text{-mod}, \quad \text{Ind}_B^A N := A \otimes_B N, \quad N \in B\text{-mod},$$

$$\text{Res}_B^A: A\text{-mod} \rightarrow B\text{-mod}, \quad \text{Res}_B^A M := \text{Hom}_A(A, M) \cong {}_B A \otimes_A M, \quad M \in A\text{-mod},$$

where ${}_B A$ denotes A considered as a (B, A) -bimodule and the left B -action on $\text{Hom}_A(A, M)$ is given by $(b, f) \mapsto f \circ {}^R b$ for $f \in \text{Hom}_A(A, M)$ and $b \in B$ (here ${}^R b$ denotes right multiplication

by b). The isomorphism above is given by the map $f \mapsto 1 \otimes f(1_A)$ for $f \in \text{Hom}_A(A, M)$. This isomorphism is natural in M and so we have an isomorphism of functors $\text{Res}_B^A \cong {}_B A \otimes -$.

3.2. Towers of algebras.

Definition 3.1 (Tower of algebras). Let $A = \bigoplus_{n \in \mathbb{N}} A_n$ be a graded algebra over a field \mathbb{F} with multiplication $\rho: A \otimes A \rightarrow A$. Then A is called a *tower of algebras* if the following conditions are satisfied:

- (TA1) Each graded piece A_n , $n \in \mathbb{N}$, is a finite dimensional algebra (with a different multiplication) with a unit 1_n . We have $A_0 \cong \mathbb{F}$.
- (TA2) The external multiplication $\rho_{m,n}: A_m \otimes A_n \rightarrow A_{m+n}$ is a homomorphism of algebras for all $m, n \in \mathbb{N}$ (sending $1_m \otimes 1_n$ to 1_{m+n}).
- (TA3) We have that A_{m+n} is a two-sided projective $(A_m \otimes A_n)$ -module with the action defined by

$$a \cdot (b \otimes c) = a\rho_{m,n}(b \otimes c) \quad \text{and} \quad (b \otimes c) \cdot a = \rho_{m,n}(b \otimes c)a,$$

for all $m, n \in \mathbb{N}$, $a \in A_{m+n}$, $b \in A_m$, $c \in A_n$.

- (TA4) For each $n \in \mathbb{N}$, the pairing (3.1) (with $B = A_n$) is perfect. (Note that this condition is automatically satisfied if \mathbb{F} is an algebraically closed field, by (3.2).)

For the remainder of this section we assume that A is a tower of algebras. We let

$$(3.3) \quad \mathcal{G}(A) = \bigoplus_{n \in \mathbb{N}} G_0(A_n) \quad \text{and} \quad \mathcal{K}(A) = \bigoplus_{n \in \mathbb{N}} K_0(A_n).$$

Then we have a perfect pairing $\langle \cdot, \cdot \rangle: \mathcal{K}(A) \times \mathcal{G}(A) \rightarrow \mathbb{Z}$ given by

$$(3.4) \quad \langle [P], [M] \rangle = \begin{cases} \dim_{\mathbb{F}}(\text{Hom}_{A_n}(P, M)) & \text{if } P \in A_n\text{-pmod and } M \in A_n\text{-mod for some } n \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

We also define a perfect pairing $\langle \cdot, \cdot \rangle: (\mathcal{K}(A) \otimes \mathcal{K}(A)) \times (\mathcal{G}(A) \otimes \mathcal{G}(A))$ by

$$\langle [P] \otimes [Q], [M] \otimes [N] \rangle = \begin{cases} \dim_{\mathbb{F}}(\text{Hom}_{A_k \otimes A_\ell}(P \otimes Q, M \otimes N)) & \text{if } P \in A_k\text{-pmod, } Q \in A_\ell\text{-pmod} \\ \quad \text{and } M \in A_k\text{-mod, } N \in A_\ell\text{-mod for some } k, \ell \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Thus we have $\langle [P] \otimes [Q], [M] \otimes [N] \rangle = \langle [P], [M] \rangle \langle [Q], [N] \rangle$.

Consider the direct sums of categories

$$A\text{-mod} := \bigoplus_{n \in \mathbb{N}} A_n\text{-mod}, \quad A\text{-pmod} := \bigoplus_{n \in \mathbb{N}} A_n\text{-pmod}.$$

For $r \in \mathbb{N}_+$, we define

$$\begin{aligned} A\text{-mod}^{\otimes r} &:= \bigoplus_{n_1, \dots, n_r \in \mathbb{N}} (A_{n_1} \otimes \cdots \otimes A_{n_r})\text{-mod}, \\ A\text{-pmod}^{\otimes r} &:= \bigoplus_{n_1, \dots, n_r \in \mathbb{N}} (A_{n_1} \otimes \cdots \otimes A_{n_r})\text{-pmod}. \end{aligned}$$

Then, for $i, j \in \{1, \dots, r\}$, $i < j$, we define $S_{ij}: A\text{-mod}^{\otimes r} \rightarrow A\text{-mod}^{\otimes r}$ to be the endofunctor that interchanges the i th and j th factors, that is, the endofunctor arising from the isomorphism

$$A_{n_1} \otimes \cdots \otimes A_{n_r} \cong A_{n_1} \otimes \cdots \otimes A_{n_{i-1}} \otimes A_{n_j} \otimes A_{n_{i+1}} \otimes \cdots \otimes A_{n_{j-1}} \otimes A_{n_i} \otimes A_{n_{j+1}} \otimes \cdots \otimes A_{n_r}.$$

We use the same notation to denote the analogous endofunctor on $A\text{-pmod}^{\otimes r}$.

We also have the following functors:

$$(3.5) \quad \begin{aligned} \nabla: A\text{-mod}^{\otimes 2} &\rightarrow A\text{-mod}, & \nabla|_{(A_m \otimes A_n)\text{-mod}} &= \text{Ind}_{A_m \otimes A_n}^{A_{m+n}}, \\ \Delta: A\text{-mod} &\rightarrow A\text{-mod}^{\otimes 2}, & \Delta|_{A_n\text{-mod}} &= \bigoplus_{k+\ell=n} \text{Res}_{A_k \otimes A_\ell}^{A_n}, \\ \eta: \text{Vect} &\rightarrow A\text{-mod}, & \eta(V) &= V \in A_0\text{-mod for } V \in \text{Vect}, \\ \varepsilon: A\text{-mod} &\rightarrow \text{Vect}, & \varepsilon(V) &= \begin{cases} V & \text{if } V \in A_0\text{-mod,} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

In the above, we have identified $A_0\text{-mod}$ with the category Vect of finite dimensional vector spaces over \mathbb{F} . Replacing $A\text{-mod}$ by $A\text{-pmod}$ above, we also have the functors ∇ , Δ , η and ε on $A\text{-pmod}$. Since the above functors are all exact (we use axiom (TA3) here), they induce a multiplication, comultiplication, unit and counit on $\mathcal{G}(A)$ and $\mathcal{K}(A)$. We use the same notation to denote these induced maps.

Since induction is always left adjoint to restriction (see, for example, [CR81, (2.19)]), ∇ is left adjoint to Δ . However, in many examples of towers of algebras (e.g. nilcoxeter algebras and 0-Hecke algebras), induction is not right adjoint to restriction. Nevertheless, we often have something quite close to this property. Any algebra automorphism ψ_n of A_n induces an isomorphism of categories $\Psi_n: A_n\text{-mod} \rightarrow A_n\text{-mod}$ (which restricts to an isomorphism of categories $\Psi_n: A_n\text{-pmod} \rightarrow A_n\text{-pmod}$) by twisting the A_n action. Then $\Psi := \bigoplus_{n \in \mathbb{N}} \Psi_n$ is an automorphism of the categories $A\text{-mod}$ and $A\text{-pmod}$. It induces automorphisms (which we also denote by Ψ) of $\mathcal{G}(A)$ and $\mathcal{K}(A)$.

Definition 3.2 (Twisted adjointness). Given a tower of algebras A , we say that induction is *twisted right adjoint* to restriction (and restriction is *twisted left adjoint* to induction) with *twisting* Ψ if there are isomorphisms of algebras $\psi_n: A_n \rightarrow A_n$, $n \in \mathbb{N}$, such that ∇ is right adjoint to $\Psi^{\otimes 2} \Delta \Psi^{-1}$.

The following lemma will be useful since many examples of towers of algebras are in fact composed of Frobenius algebras. We refer the reader to [SY11] for background on Frobenius algebras.

Lemma 3.3. *If each A_n , $n \in \mathbb{N}$, is a Frobenius algebra, then induction is twisted right adjoint to restriction, with twisting given by ψ_n being the inverse of the Nakayama automorphism of A_n .*

Proof. This follows from [Kho01, Lem. 1] by taking $B_1 = A_{m+n}$, $B_2 = A_m \otimes A_n$ and N to be A_{m+n} , considered as an $(A_{m+n}, A_m \otimes A_n)$ -bimodule in the natural way. \square

Definition 3.4 (Strong tower of algebras). We say that a tower of algebras A is *strong* if induction is twisted right adjoint to restriction and we have an isomorphism of functors

$$(3.6) \quad \Delta \nabla \cong \nabla^{\otimes 2} S_{23} \Delta^{\otimes 2}.$$

Remark 3.5. The isomorphism (3.6) is a compatibility between induction and restriction that is analogous to the well known Mackey theory for finite groups. It implies that $\mathcal{K}(A)$ and $\mathcal{G}(A)$ are Hopf algebras under the operations defined above (see [BL09, Th. 3.5] – although the authors of that paper work over \mathbb{C} and they assume that the external multiplication maps $\rho_{m,n}$ are injective, the arguments hold in the more general setting considered here).

Definition 3.6 (Dualizing tower of algebras). We say that a tower of algebras A is *dualizing* if, under the operations defined above, $\mathcal{K}(A)$ and $\mathcal{G}(A)$ are Hopf algebras which are dual, in the sense of Definition 2.4 (with $\mathbb{k} = \mathbb{Z}$), under the bilinear form (3.4) (i.e. the perfect pairing (3.4) is a Hopf pairing).

Proposition 3.7. *Suppose A is a strong tower of algebras with twisting Ψ . Then the following statements are equivalent.*

- (a) *The tower A is dualizing.*
- (b) *We have $\Psi^{\otimes 2} \Delta \Psi^{-1}(P) \cong \Delta(P)$ for all $P \in A\text{-pmod}$.*
- (c) *We have $\Psi^{\otimes 2} \Delta \Psi^{-1} = \Delta$ as endomorphisms of $\mathcal{K}(A)$.*

In particular A is dualizing if each A_n is a symmetric algebra (i.e. if $\Psi = \text{Id}$) or, more generally, if Ψ acts trivially on $\mathcal{K}(A)$.

Proof. First assume that (b) holds. With one exception, the proof that A is dualizing then proceeds exactly as in the proof of [BL09, Th. 3.6] since (3.6) implies axiom (5) in [BL09, §3.1]. (Although the authors of that paper work over \mathbb{C} and they assume that the external multiplication maps $\rho_{m,n}$ are injective, the arguments hold in the more general setting considered here.) The one exception is in the proof that

$$(3.7) \quad \langle \Delta([P]), [M] \otimes [N] \rangle = \langle [P], \nabla([M] \otimes [N]) \rangle,$$

for all $M \in A_m\text{-mod}$, $N \in A_n\text{-mod}$, $P \in A_{m+n}\text{-pmod}$. However, under our assumptions, we have

$$\text{Hom}_{A_{m+n}}(P, \nabla(M \otimes N)) \cong \text{Hom}_{A_m \otimes A_n}(\Psi^{\otimes 2} \Delta \Psi^{-1}(P), M \otimes N) \cong \text{Hom}_{A_m \otimes A_n}(\Delta(P), M \otimes N),$$

which immediately implies (3.7). Thus A is dualizing.

Now suppose A is dualizing. Then, for all $P \in A\text{-pmod}$ and $M \in A\text{-mod}^{\otimes 2}$, we have

$$\langle \Psi^{\otimes 2} \Delta \Psi^{-1}([P]), [M] \rangle = \langle [P], \nabla([M]) \rangle = \langle \Delta([P]), [M] \rangle,$$

where the first equality holds by the assumption that induction is twisted right adjoint to restriction and the second equality holds by the assumption that the tower is dualizing. Then (c) follows from the nondegeneracy of the bilinear form.

The fact that (b) and (c) are equivalent follows from the fact that every short exact sequence of projective modules splits. Thus, for $P, Q \in A\text{-pmod}$, we have $[P] = [Q]$ in $\mathcal{K}(A)$ if and only if $P \cong Q$. \square

Remark 3.8. It is crucial in (b) that P be a projective module. The isomorphism does not hold, in general, for arbitrary modules, even if the tower is dualizing. For instance, the tower of 0-Hecke algebras is dualizing (see Corollary 4.6), but one can show that $\Psi^{\otimes 2} \Delta \Psi^{-1} \cong S_{12} \Delta$ (see Lemma 4.4). Then (b) corresponds to the fact that the comultiplication on NSym is cocommutative. However, the comultiplication on QSym is not cocommutative and thus (b) does not hold, in general, if P is not projective. We refer the reader to Section 8 for further details on the tower of 0-Hecke algebras.

3.3. The Heisenberg double associated to a tower of algebras. In this section we apply the constructions of Section 2 to the dual pair $(\mathcal{G}(A), \mathcal{K}(A))$ arising from a dualizing tower of algebras A . We also see that some natural subalgebras of the Heisenberg double arise in this situation.

Definition 3.9 ($\mathfrak{h}(A)$, $\mathcal{F}(A)$, $\mathcal{G}_{\text{proj}}(A)$). Suppose A is a dualizing tower of algebras. We let $\mathfrak{h}(A) = \mathfrak{h}(\mathcal{G}(A), \mathcal{K}(A))$ and $\mathcal{F}(A) = \mathcal{F}(\mathcal{G}(A), \mathcal{K}(A))$. For each $n \in \mathbb{N}$, $A_n\text{-pmod}$ is a full subcategory of $A_n\text{-mod}$. The inclusion functor induces the *Cartan map* $\mathcal{K}(A) \rightarrow \mathcal{G}(A)$. Let $\mathcal{G}_{\text{proj}}(A)$ denote the image of the Cartan map.

For the remainder of this section, we fix a dualizing tower of algebras A and let

$$(3.8) \quad H^- = \mathcal{K}(A), \quad H^+ = \mathcal{G}(A), \quad H_{\text{proj}}^+ = \mathcal{G}_{\text{proj}}(A), \quad \mathfrak{h} = \mathfrak{h}(A), \quad \mathcal{F} = \mathcal{F}(A).$$

To avoid confusion between $\mathcal{G}(A)$ and $\mathcal{K}(A)$, we will write $[M]_+$ to denote the class of a finitely generated (possibly projective) A_n -module in H^+ and $[M]_-$ to denote the class of a finitely generated projective A_n -module in H^- . If $P \in A_p\text{-pmod}$ and $N \in (A_{n-p} \otimes A_p)\text{-mod}$, then we have a natural A_{n-p} -module structure on $\text{Hom}_{A_p}(P, N)$ given by

$$(3.9) \quad (a \cdot f)(b) = (a \otimes 1) \cdot (f(b)), \quad a \in A_{n-m}, \quad f \in \text{Hom}_{A_p}(P, N), \quad b \in P.$$

Lemma 3.10. *If $p, n \in \mathbb{N}$, $P \in A_p\text{-pmod}$ and $N \in A_n\text{-mod}$, then we have*

$$[P]_- \cdot [N]_+ = \begin{cases} 0 & \text{if } p > n, \\ [\text{Hom}_{A_p}(P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N)]_+ & \text{if } p \leq n. \end{cases}$$

Here \cdot denotes the action of \mathfrak{h} on \mathcal{F} and $\text{Hom}_{A_p}(P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N)$ is viewed as an A_{n-p} -module as in (3.9).

Proof. The case $p > n$ follows immediately from the fact that $H_{n-p}^+ = 0$ if $p > n$. Assume $p \leq n$. For $R \in A_{n-p}\text{-pmod}$, we have

$$\begin{aligned} \langle [R]_-, [P]_- \cdot [N]_+ \rangle &= \langle [R]_- [P]_-, [N]_+ \rangle \\ &= \langle \nabla([R]_- \otimes [P]_-), [N]_+ \rangle \\ &= \langle [R]_- \otimes [P]_-, \Delta([N]_+) \rangle \\ &= \dim_{\mathbb{F}} \text{Hom}_{A_{n-p} \otimes A_p}(R \otimes P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N) \\ &= \dim_{\mathbb{F}} \text{Hom}_{A_{n-p}}(R, \text{Hom}_{A_p}(P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N)) \\ &= \langle [R]_-, [\text{Hom}_{A_p}(P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N)]_+ \rangle. \end{aligned}$$

The result then follows from the nondegeneracy of the bilinear form. \square

Lemma 3.11. *Suppose \mathbb{k} is a commutative ring and R and S are \mathbb{k} -algebras. Furthermore, suppose that P is a projective S -module and Q is a projective $(R \otimes S)$ -module. Then $\text{Hom}_S(P, Q)$ is a projective R -module.*

Proof. If P is a projective S -module and Q is a projective $(R \otimes S)$ -module then there exist $s, t \in \mathbb{N}$, an S -module P' and an $(R \otimes S)$ -module Q' such that $P \oplus P' \cong S^s$ and $Q \oplus Q' \cong (R \otimes S)^t$. Then we have

$$\begin{aligned} \text{Hom}_S(P, Q) \oplus \text{Hom}_S(P, Q') \oplus \text{Hom}_S(P', (R \otimes S)^t) \\ \cong \text{Hom}_S(S^s, (R \otimes S)^t) \cong \text{Hom}_S(S, (R \otimes S)^{st}) \cong R^{st}. \end{aligned}$$

Thus $\text{Hom}_S(P, Q)$ is a projective R -module. \square

Proposition 3.12. *We have that H_{proj}^+ is a subalgebra of H^+ that is invariant under the left regular action of H^- .*

Proof. Assume $P \in A_p\text{-pmod}$ and $Q \in A_q\text{-pmod}$. As in the proof of [BL09, Prop. 3.2], we have that $\text{Ind}_{A_p \otimes A_q}^{A_{p+q}}(P \otimes Q)$ is a projective A_{p+q} -module. It follows that H_{proj}^+ is a subalgebra of H^+ .

By Lemma 3.10, it remains to show that

$$\text{Hom}_{A_p}(P, \text{Res}_{A_{q-p} \otimes A_p}^{A_q} Q) \in A_{q-p}\text{-pmod}.$$

Again, as in the proof of [BL09, Prop. 3.2], we have that $\text{Res}_{A_{q-p} \otimes A_p}^{A_q} Q$ is a projective $(A_{q-p} \otimes A_p)$ -module. The result then follows from Lemma 3.11. \square

Definition 3.13 (The projective Heisenberg double $\mathfrak{h}_{\text{proj}}(A)$). By Proposition 3.12, $\mathfrak{h}_{\text{proj}} = \mathfrak{h}_{\text{proj}}(A) := H_{\text{proj}}^+ \# H^-$ is a subalgebra of \mathfrak{h} . In other words, $\mathfrak{h}_{\text{proj}}$ is the subalgebra of \mathfrak{h} generated by H_{proj}^+ and H^- (viewing the latter two as \mathbb{Z} -submodules of \mathfrak{h} as in Definition 2.6). We call $\mathfrak{h}_{\text{proj}}$ the *projective Heisenberg double* associated to A .

Definition 3.14 (Fock space $\mathcal{F}_{\text{proj}}(A)$ of $\mathfrak{h}_{\text{proj}}$). By Proposition 3.12, the algebra $\mathfrak{h}_{\text{proj}}$ acts on H_{proj}^+ . We call this the *lowest weight Fock space representation* of $\mathfrak{h}_{\text{proj}}$ and denote it by $\mathcal{F}_{\text{proj}} = \mathcal{F}_{\text{proj}}(A)$. Note that this representation is generated by the lowest weight vacuum vector $1 \in H_{\text{proj}}^+$.

Proposition 3.15. *The Fock space $\mathcal{F}_{\text{proj}}$ of $\mathfrak{h}_{\text{proj}}$ has the following properties.*

- (a) *The only submodules of $\mathcal{F}_{\text{proj}}$ are those submodules of the form $n\mathcal{F}_{\text{proj}}$ for $n \in \mathbb{Z}$.*
- (b) *Let $\mathbb{Z}^- \cong \mathbb{Z}$ (isomorphism of \mathbb{Z} -modules) be the representation of H^- such that H_n^- acts as zero for all $n > 0$ and $H_0^- \cong \mathbb{Z}$ acts by left multiplication. Then $\mathcal{F}_{\text{proj}}$ is isomorphic to the induced module $\text{Ind}_{H^-}^{\mathfrak{h}_{\text{proj}}} \mathbb{Z}^-$ as an $\mathfrak{h}_{\text{proj}}$ -module.*
- (c) *Any representation of $\mathfrak{h}_{\text{proj}}$ generated by a lowest weight vacuum vector is isomorphic to $\mathcal{F}_{\text{proj}}$.*

Proof. This is an immediate consequence of Theorem 2.11, taking $X^+ = H_{\text{proj}}^+$. \square

Note that the lowest weight Fock space $\mathcal{F}_{\text{proj}}$ is not a faithful $\mathfrak{h}_{\text{proj}}$ -module in general (see Section 8.5), in contrast to the case for \mathfrak{h} (see Theorem 2.11(d)). However, we can define a highest weight Fock space of $\mathfrak{h}_{\text{proj}}$ that is faithful. Consider the augmentation algebra homomorphism $\epsilon^+ : H_{\text{proj}}^+ \rightarrow \mathbb{Z}$ uniquely determined by $\epsilon^+(H_n^+ \cap H_{\text{proj}}^+) = 0$ for $n > 0$. Let \mathbb{Z}_ϵ^+ denote the corresponding H_{proj}^+ -module. We call the induced module $\mathcal{F}_{\text{proj}}^- := \text{Ind}_{H_{\text{proj}}^+}^{\mathfrak{h}_{\text{proj}}} \mathbb{Z}_\epsilon^+$ the *highest weight Fock space representation* of $\mathfrak{h}_{\text{proj}}$. It is generated by the highest weight vacuum vector $1 \in \mathbb{Z}_\epsilon^+$.

Proposition 3.16. *The highest weight Fock space representation $\mathcal{F}_{\text{proj}}^-$ of $\mathfrak{h}_{\text{proj}}$ is faithful.*

Proof. We have $\mathfrak{h}_{\text{proj}} \cong H_{\text{proj}}^+ \otimes H^-$ as \mathbb{k} -modules and so $\mathcal{F}_{\text{proj}}^- \cong H^-$ as \mathbb{k} -modules. The action of $\mathfrak{h}_{\text{proj}}$ on $\mathcal{F}_{\text{proj}}^-$ is simply the restriction of the natural action of \mathfrak{h} on H^- , which is faithful by (an obvious highest weight analogue of) Theorem 2.11(d). \square

3.4. Categorification of Fock space. In this section we prove our main result, the categorification of the Fock space representation of the Heisenberg double. We continue to fix a dualizing tower of algebras A and to use the notation of (3.8).

Recall the direct sums of categories

$$A\text{-mod} = \bigoplus_{n \in \mathbb{N}} A_n\text{-mod}, \quad A\text{-pmod} = \bigoplus_{n \in \mathbb{N}} A_n\text{-pmod}.$$

For each $M \in A_m\text{-mod}$, $m \in \mathbb{N}$, define the functor $\text{Ind}_M: A\text{-mod} \rightarrow A\text{-mod}$ by

$$\text{Ind}_M(N) = \text{Ind}_{A_m \otimes A_n}^{A_{m+n}}(M \otimes N) \in A_{m+n}\text{-mod}, \quad N \in A_n\text{-mod}, \quad n \in \mathbb{N}.$$

For each $P \in A_p\text{-pmod}$, $p \in \mathbb{N}$, define the functor $\text{Res}_P: A\text{-mod} \rightarrow A\text{-mod}$ by

$$\text{Res}_P(N) = \text{Hom}_{A_p}(P, \text{Res}_{A_{n-p} \otimes A_p}^{A_n} N) \in A_{n-p}\text{-mod}, \quad N \in A_n\text{-mod}, \quad n \in \mathbb{N},$$

where $\text{Res}_P(N)$ is interpreted to be the zero object of $A\text{-mod}$ if $n - p < 0$.

As in the proof of Proposition 3.12, we see that

$$\text{Ind}_P(A\text{-pmod}) \subseteq A\text{-pmod}, \quad \text{Res}_P(A\text{-pmod}) \subseteq A\text{-pmod} \quad \text{for all } P \in A\text{-pmod}.$$

Thus we have the induced functors $\text{Ind}_P, \text{Res}_P: A\text{-pmod} \rightarrow A\text{-pmod}$ for $P \in A\text{-pmod}$.

Since the functors Ind_M and Res_P are exact for all $M \in A\text{-mod}$ and $P \in A\text{-pmod}$, they induce endomorphisms $[\text{Ind}_M]$ and $[\text{Res}_P]$ of $\mathcal{G}(A)$. Similarly, Ind_P and Res_P induce endomorphisms $[\text{Ind}_P]$ and $[\text{Res}_P]$ of $\mathcal{G}_{\text{proj}}(A)$ for all $P \in A\text{-pmod}$.

Proposition 3.17. *Suppose A is a dualizing tower of algebras.*

(a) *For all $M, N \in A\text{-mod}$ and $P \in A\text{-pmod}$, we have*

$$([M] \# [P])([N]) = [\text{Ind}_M] \circ [\text{Res}_P]([N]) = [\text{Ind}_M \circ \text{Res}_P(N)] \in \mathcal{G}(A).$$

(b) *For all $Q, P, R \in A\text{-pmod}$, we have*

$$([Q] \# [P])([R]) = [\text{Ind}_Q] \circ [\text{Res}_P]([R]) = [\text{Ind}_Q \circ \text{Res}_P(R)] \in \mathcal{G}_{\text{proj}}(A).$$

Proof. This follows from the definition of the multiplication in $\mathcal{G}(A)$ and Lemma 3.10. \square

Part (a) (resp. part (b)) of Proposition 3.17 shows how the action of \mathfrak{h} on \mathcal{F} (resp. $\mathfrak{h}_{\text{proj}}$ on $\mathcal{F}_{\text{proj}}$) is induced by functors on $\bigoplus_{n \geq 0} A_n\text{-mod}$ (resp. $\bigoplus_{n \geq 0} A_n\text{-pmod}$). Typically a *categorification* of a representation consists of isomorphisms of such functors which lift the algebra relations. As we now describe, this can be done if the tower of algebras is strong.

First note that the algebra structure on \mathfrak{h} is uniquely determined by the fact that H^\pm are subalgebras and by the relation

$$(3.10) \quad xa = \sum_{(x)} {}^R x_{(1)} {}^*(a) x_{(2)}, \quad x \in H^-, \quad a \in H^+,$$

between H^- and H^+ . Since the natural action of \mathfrak{h} on H^+ is faithful by Theorem 2.11(d), equation (3.10) is equivalent to the equalities in $\text{End } H^+$

$$(3.11) \quad {}^R x^* \circ {}^L a = \nabla \left({}^R \Delta(x) {}^*(a \otimes -) \right), \quad x \in H^-, \quad a \in H^+.$$

For $Q \in (A_p \otimes A_q)\text{-pmod}$, $M \in A_m\text{-mod}$, and $N \in A_n\text{-mod}$, define

$$\text{Res}_Q(M \otimes N) := \text{Hom}_{A_p \otimes A_q} \left(Q, S_{23} \left(\text{Res}_{A_{m-p} \otimes A_p}^{A_m} M \otimes \text{Res}_{A_{n-q} \otimes A_q}^{A_n} N \right) \right).$$

For $Q \in A\text{-mod}^{\otimes 2}$, we let Res_Q denote the corresponding sum of functors.

Theorem 3.18. *Suppose that A is a strong tower of algebras. Then we have the following isomorphisms of functors for all $M, N \in A\text{-mod}$ and $P, Q \in A\text{-pmod}$.*

$$(3.12) \quad \text{Ind}_M \circ \text{Ind}_N \cong \text{Ind}_{\nabla(M \otimes N)},$$

$$(3.13) \quad \text{Res}_P \circ \text{Res}_Q \cong \text{Res}_{\nabla(P \otimes Q)},$$

$$(3.14) \quad \text{Res}_P \circ \text{Ind}_M \cong \nabla \text{Res}_{\Psi^{\otimes 2} \Delta \Psi^{-1}(P)}(M \otimes -).$$

In particular, if A is dualizing, then the above yields a categorification of the lowest weight Fock space representations of $\mathfrak{h}(A)$ and $\mathfrak{h}_{\text{proj}}(A)$.

The isomorphisms (3.12) and (3.13) categorify the multiplication in $\mathcal{G}(A)$ and $\mathcal{K}(A)$ respectively. If A is dualizing, then, in light of Proposition 3.7, the isomorphism (3.14) categorifies the relation (3.11). Thus, Theorem 3.18 provides a categorification of the lowest weight Fock space representation $\mathcal{F}(A)$ of $\mathfrak{h}(A)$. If we restrict the induction and restriction functors to $A\text{-pmod}$ and require $M, N \in A\text{-pmod}$ in the statement of the theorem, then we obtain a categorification of the lowest weight Fock space representation $\mathcal{F}_{\text{proj}}(A)$ of $\mathfrak{h}_{\text{proj}}(A)$. Note that the categorification in Theorem 3.18 does not rely on a particular presentation of the Heisenberg double $\mathfrak{h}(A)$ (see Remark 6.1), in contrast to many other categorification statements appearing in the literature.

Proof of Theorem 3.18. Suppose $M \in A_m\text{-mod}$, $N \in A_n\text{-mod}$, $P \in A_p\text{-pmod}$, $Q \in A_q\text{-pmod}$, and $L \in A_\ell\text{-mod}$. Then we have

$$\begin{aligned} \text{Ind}_M \circ \text{Ind}_N(L) &= \text{Ind}_{A_m \otimes A_{n+\ell}}^{A_{m+n+\ell}} \left(M \otimes \text{Ind}_{A_n \otimes A_\ell}^{A_{n+\ell}}(N \otimes L) \right) \\ &\cong \text{Ind}_{A_m \otimes A_{n+\ell}}^{A_{m+n+\ell}} \text{Ind}_{A_m \otimes A_n \otimes A_\ell}^{A_m \otimes A_{n+\ell}}(M \otimes N \otimes L) \\ &\cong \text{Ind}_{A_m \otimes A_n \otimes A_\ell}^{A_{m+n+\ell}}(M \otimes N \otimes L) \\ &\cong \text{Ind}_{A_{m+n} \otimes A_\ell}^{A_{m+n+\ell}} \text{Ind}_{A_m \otimes A_n \otimes A_\ell}^{A_{m+n} \otimes A_\ell}(M \otimes N \otimes L) \\ &\cong \text{Ind}_{A_{m+n} \otimes A_\ell}^{A_{m+n+\ell}} \left(\text{Ind}_{A_m \otimes A_n}^{A_{m+n}}(M \otimes N) \otimes L \right) \\ &\cong \text{Ind}_{\nabla(M \otimes N)} L. \end{aligned}$$

Since each of the above isomorphisms is natural in L , this proves (3.12).

Similarly, we have

$$\begin{aligned} \text{Res}_P \circ \text{Res}_Q(L) &= \text{Hom}_{A_p}(P, \text{Res}_{A_{\ell-q-p} \otimes A_p}^{A_{\ell-q}} \text{Hom}_{A_q}(Q, \text{Res}_{A_{\ell-q} \otimes A_q}^{A_\ell} L)) \\ &\cong \text{Hom}_{A_p}(P, \text{Hom}_{A_q}(Q, \text{Res}_{A_{\ell-p-q} \otimes A_p \otimes A_q}^{A_\ell} L)) \\ &\cong \text{Hom}_{A_p \otimes A_q}(P \otimes Q, \text{Res}_{A_{\ell-p-q} \otimes A_p \otimes A_q}^{A_\ell} L) \\ &\cong \text{Hom}_{A_p \otimes A_q}(P \otimes Q, \text{Res}_{A_{\ell-p-q} \otimes A_p \otimes A_q}^{A_{\ell-p-q} \otimes A_{p+q}} \text{Res}_{A_{\ell-p-q} \otimes A_{p+q}}^{A_\ell} L) \\ &\cong \text{Hom}_{A_{p+q}}(\text{Ind}_{A_p \otimes A_q}^{A_{p+q}}(P \otimes Q), \text{Res}_{A_{\ell-p-q} \otimes A_{p+q}}^{A_\ell} L) \\ &\cong \text{Res}_{\nabla(P \otimes Q)} L. \end{aligned}$$

Since each of the above isomorphisms is natural in L , this proves (3.13).

Finally, we have

$$\text{Res}_P \circ \text{Ind}_M(L) = \text{Hom}_{A_p}(P, \text{Res}_{A_{m+\ell-p} \otimes A_p}^{A_{m+\ell}} \text{Ind}_{A_m \otimes A_\ell}^{A_{m+\ell}}(M \otimes L))$$

$$\begin{aligned}
&\cong \mathrm{Hom}_{A_p}(P, \bigoplus_{s+t=p} \mathrm{Ind}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_{m+\ell-p} \otimes A_p} \mathrm{Res}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_m \otimes A_\ell}(M \otimes L)) \\
&\cong \mathrm{Hom}_{A_p}(P, \bigoplus_{s+t=p} \mathrm{Ind}_{A_{m-s} \otimes A_{\ell-t} \otimes A_p}^{A_{m+\ell-p} \otimes A_p} \mathrm{Ind}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_{m-s} \otimes A_{\ell-t} \otimes A_p} \mathrm{Res}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_m \otimes A_\ell}(M \otimes L)) \\
&\cong \bigoplus_{s+t=p} \mathrm{Ind}_{A_{m-s} \otimes A_{\ell-t}}^{A_{m+\ell-p}} \mathrm{Hom}_{A_p}(P, \mathrm{Ind}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_{m-s} \otimes A_{\ell-t} \otimes A_p} \mathrm{Res}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_m \otimes A_\ell}(M \otimes L)) \\
&\cong \bigoplus_{s+t=p} \mathrm{Ind}_{A_{m-s} \otimes A_{\ell-t}}^{A_{m+\ell-p}} \mathrm{Hom}_{A_s \otimes A_t}(\Psi^{\otimes 2} \mathrm{Res}_{A_s \otimes A_t}^{A_p} \Psi^{-1}(P), \mathrm{Res}_{A_{m-s} \otimes A_s \otimes A_{\ell-t} \otimes A_t}^{A_m \otimes A_\ell}(M \otimes L)) \\
&\cong \nabla \mathrm{Res}_{\Psi^{\otimes 2} \Delta \Psi^{-1}(P)}(M \otimes L),
\end{aligned}$$

where the first isomorphism follows from (3.6). Since all of the above isomorphisms are natural in L , this proves (3.14).

The final assertion of the theorem follows from Proposition 3.7 as explained in the paragraph following the statement of the theorem. \square

4. HECKE-LIKE TOWERS OF ALGEBRAS

In the remainder of the paper we will be studying well known examples of towers of algebras. These examples are all quotients of groups algebras of braid groups by quadratic relations. In this subsection, we prove that all such towers of algebras are strong and dualizing. In this section, \mathbb{F} is an algebraically closed field unless otherwise specified.

Definition 4.1 (Hecke-like algebras and towers). We say that the \mathbb{F} -algebra B is *Hecke-like* (of degree n) if it is generated by elements T_1, \dots, T_{n-1} , subject to the relations

$$\begin{aligned}
T_i T_j &= T_j T_i, & |i - j| > 1, \\
T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, & i = 1, \dots, n-2, \\
T_i^2 &= c T_i + d, & i = 1, \dots, n-1,
\end{aligned}$$

for some $c, d \in \mathbb{F}$ (independent of i). In other words, B is Hecke-like if it is a quotient of the group algebra of the braid group by quadratic relations (the last set of relations above).

If, for $n \in \mathbb{N}$, the algebra A_n is a Hecke-like algebra of degree n , and the constants c, d above are independent of n , then we can define an external multiplication on $A = \bigoplus_{n \in \mathbb{N}} A_n$ by

$$(4.1) \quad \rho_{m,n}: A_m \otimes A_n \rightarrow A_{m+n}, \quad T_i \otimes 1 \mapsto T_i, \quad 1 \otimes T_i \mapsto T_{m+i}.$$

Axioms (TA1) and (TA2) follow immediately. Furthermore, it follows from Lemma 4.2 below that, as a left $(A_m \otimes A_n)$ -module, we have

$$A_{m+n} = \bigoplus_{w \in X_{m,n}} (A_m \otimes A_n) T_w,$$

where $X_{m,n}$ is a set of minimal length representatives of the cosets $(A_m \otimes A_n) \backslash A_{m+n}$. Thus A_{m+n} is a projective left $(A_m \otimes A_n)$ -module. Similarly, it is also a projective right $(A_m \otimes A_n)$ -module and so axiom (TA3) is satisfied. Finally (TA4) is satisfied since \mathbb{F} is algebraically closed. We call the resulting tower of algebras a *Hecke-like tower of algebras*.

Lemma 4.2. *Suppose that B is a Hecke-like algebra of degree n .*

- (a) *If, for $w \in S_n$, we define $T_w = T_{i_1} \cdots T_{i_r}$, where $s_{i_1} \cdots s_{i_r}$ is a reduced decomposition of w (these elements are well defined by the braid relations), then $\{T_w \mid w \in S_n\}$ is a basis of B . In particular, the dimension of B is $n!$.*

(b) We have

$$T_{w_1}T_{w_2} = T_{w_1w_2} \quad \text{for all } w_1, w_2 \in S_n \text{ such that } \ell(w_1w_2) = \ell(w_1) + \ell(w_2),$$

where ℓ is the usual length function on S_n .

(c) The algebra B is a Frobenius algebra with trace map $\lambda: B \rightarrow \mathbb{F}$ given by $\lambda(T_w) = \delta_{w, w_0}$, where w_0 is the longest element of S_n . The corresponding Nakayama automorphism is the map $\psi_n: B \rightarrow B$ given by $\psi_n(T_i) = T_{n-i}$.

Proof. The proof of parts (a) and (b) is analogous to the proof for the usual Hecke algebra of type A and is left to the reader. It remains to prove part (c).

Suppose B is a Hecke-like algebra of degree n . To show that B is a Frobenius algebra with trace map λ , it suffices to show that $\ker \lambda$ contains no nonzero left ideals. Let I be a nonzero ideal of B . Then choose a nonzero element $b = \sum_{w \in S_n} a_w T_w \in I$ and let τ be a maximal length element of the set $\{w \in S_n \mid a_w \neq 0\}$. Then we have $\lambda(bT_{\tau^{-1}w_0}) = a_\tau \neq 0$. Thus I is not contained in $\ker \lambda$.

To show that ψ_n is the Nakayama automorphism, it suffices to show that $\lambda(T_w T_i) = \lambda(T_{n-i} T_w)$ for all $i \in \{1, \dots, n-1\}$ and $w \in S_n$. We break the proof into four cases.

Case 1: $\ell(w) \leq \ell(w_0) - 2$. In this case we clearly have $\lambda(T_w T_i) = 0 = \lambda(T_{n-i} T_w)$.

Case 2: $w = w_0$. Then we can write $w = w_0 = \tau s_i$ for some $\tau \in S_n$ with $\ell(\tau) = \ell(w_0) - 1$. Then $\lambda(T_w T_i) = \lambda(cT_{w_0} + dT_\tau) = c$. Now, since $w_0 s_i = s_{n-i} w_0$, we have $w = w_0 = s_{n-i} \tau$. Thus $\lambda(T_{n-i} T_w) = \lambda(cT_{w_0} + dT_\tau) = c = \lambda(T_w T_i)$.

Case 3: $ws_i = w_0$. Since, as noted above, we have $w_0 s_i = s_{n-i} w_0$, it follows that $s_{n-i} w = w_0$ and so $\lambda(T_w T_i) = \lambda(T_{w_0}) = \lambda(T_{n-i} T_w)$.

Case 4: $\ell(w) = \ell(w_0) - 1$, but $ws_i \neq w_0$. Then we have $w = \tau s_i$ for some $\tau \in S_n$ with $\ell(\tau) = \ell(w) - 1$. Thus $\lambda(T_w T_i) = \lambda(cT_w + dT_\tau) = 0$. Using again the equality $w_0 s_i = s_{n-i} w_0$, we have $s_{n-i} w \neq w_0$. Then an analogous argument shows that $\lambda(T_{n-i} T_w) = 0$. \square

When considering a Hecke-like tower of algebras, we will always assume that the twisting Ψ is given by the Nakayama automorphisms ψ_n described above (see Proposition 3.3).

Proposition 4.3. *All Hecke-like towers of algebras are strong.*

Proof. Suppose $A = \bigoplus_{n \in \mathbb{N}} A_n$ is a Hecke-like tower of algebras. We formulate the isomorphism (3.6) in terms of bimodules. Fix n, m, k, ℓ such that $n + m = k + \ell$ and set $N = n + m$. Let ${}_{(k, \ell)}(A_N)_{(n, m)}$ denote A_N , thought of as a $(A_k \otimes A_\ell, A_n \otimes A_m)$ -bimodule in the natural way. Then we have

$$\text{Res}_{A_k \otimes A_\ell}^{A_N} \text{Ind}_{A_n \otimes A_m}^{A_N} \cong {}_{(k, \ell)}(A_N)_{(n, m)} \otimes -.$$

On the other hand, for each r satisfying $k - m = n - \ell \leq r \leq \min\{n, k\}$, we have

$$\begin{aligned} \text{Ind}_{A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell+r-n}}^{A_k \otimes A_\ell} \text{Res}_{A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell+r-n}}^{A_n \otimes A_m} &\cong B_r \otimes -, \quad \text{where} \\ B_r &= (A_k \otimes A_\ell) \otimes_{A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell+r-n}} (A_n \otimes A_m), \end{aligned}$$

and where we view $A_k \otimes A_\ell$ as a right $(A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell+r-n})$ -module via the map $a_1 \otimes a_2 \otimes a_3 \otimes a_4 \mapsto a_1 a_3 \otimes a_2 a_4$. (This corresponds to the functor S_{23} appearing in (3.6).) Therefore, in order to prove the isomorphism (3.6), it suffices to prove that we have an isomorphism of bimodules

$${}_{(k, \ell)}(A_N)_{(n, m)} \cong \bigoplus_{r=n-\ell}^{\min\{n, k\}} B_r.$$

Now, we have one double coset in $S_k \times S_\ell \setminus S_N/S_n \times S_m$ for each r satisfying $k - m = n - \ell \leq r \leq \min\{n, k\}$ (see, for example, [Zel81, Appendix 3, p. 170]). Precisely, the double coset C_r corresponding to r consists of the permutations $w \in S_n$ satisfying

$$\begin{aligned} |w(\{1, \dots, n\}) \cap \{1, 2, \dots, k\}| &= r, \\ |w(\{n+1, \dots, N\}) \cap \{1, 2, \dots, k\}| &= k - r, \\ |w(\{1, \dots, n\}) \cap \{k+1, \dots, N\}| &= n - r, \\ |w(\{n+1, \dots, N\}) \cap \{k+1, \dots, N\}| &= \ell - n + r = m - k + r. \end{aligned}$$

Thus the cardinality of the double coset C_r is

$$|C_r| = \frac{k! \ell! m! n!}{r! (k-r)! (n-r)! (\ell-n+r)!}.$$

The permutation $w_r \in S_N$ given by

$$w_r(i) = \begin{cases} i & \text{if } 1 \leq i \leq r, \\ i - r + k & \text{if } r < i \leq n, \\ i - n + r & \text{if } n < i \leq n + k - r, \\ i & \text{if } n + k - r < i \leq N, \end{cases}$$

is a minimal length representative of C_r . It then follows from Lemma 4.2(b) that

$$\begin{aligned} T_{w_r} T_i &= T_i T_{w_r} & \text{if } 1 \leq i < r \text{ or } n + k - r < i < N, \\ T_{w_r} T_i &= T_{i-r+k} T_{w_r} & \text{if } r < i < n, \\ T_{w_r} T_i &= T_{i-n+r} T_{w_r} & \text{if } n < i < n + k - r. \end{aligned}$$

Thus,

$$(4.2) \quad T_{w_r}(a_1 \otimes a_2 \otimes a_3 \otimes a_4) = (a_1 \otimes a_3 \otimes a_2 \otimes a_4) T_{w_r}$$

for all $a_1 \in A_r$, $a_2 \in A_{n-r}$, $a_3 \in A_{k-r}$, $a_4 \in A_{\ell+r-n}$.

Define $B'_r \subseteq {}_{(k,\ell)}(A_N)_{(n,m)}$ to be the sub-bimodule generated by T_{w_r} . It follows from Lemma 4.2 that ${}_{(k,\ell)}(A_N)_{(n,m)} = \bigoplus_{r=n-\ell}^{\min\{n,k\}} B'_r$ and that $\dim_{\mathbb{F}} B'_r = |C_r|$. It also follows that the dimension of $A_k \otimes A_\ell$ as a right module over $A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell-n+r}$ is $k! \ell! / r! (n-r)! (k-r)! (\ell-n+r)!$ and that the dimension of $A_m \otimes A_n$ as a left module over $A_r \otimes A_{n-r} \otimes A_{k-r} \otimes A_{\ell+r-n}$ is $m! n! / r! (n-r)! (k-r)! (\ell-n+r)!$. Therefore, $\dim_{\mathbb{F}} B_r = |C_r| = \dim_{\mathbb{F}} B'_r$. Now consider the $(A_k \otimes A_\ell, A_n \otimes A_m)$ -bimodule map $B_r \rightarrow B'_r$ uniquely determined by

$$1_{A_k \otimes A_\ell} \otimes 1_{A_n \otimes A_m} \mapsto T_{w_r},$$

which is well defined by (4.2). This map is surjective, and thus is an isomorphism by dimension considerations. \square

Lemma 4.4. *If A is a Hecke-like tower of algebras, then we have an isomorphism of functors $\Psi^{\otimes 2} \Delta \Psi^{-1} \cong S_{12} \Delta$ on A -mod (hence also on A -pmod).*

Proof. It suffices to prove that, for $m, n \in \mathbb{N}$, we have an isomorphism of functors

$$(\Psi_m \otimes \Psi_n) \circ \text{Res}_{A_m \otimes A_n}^{A_{m+n}} \circ \Psi_{m+n}^{-1} \cong S_{12} \circ \text{Res}_{A_n \otimes A_m}^{A_{m+n}}.$$

Describing each functor above as tensoring on the left by the appropriate bimodule, it suffices to prove that we have an isomorphism of bimodules

$$(4.3) \quad (A_m^\psi \otimes A_n^\psi) \otimes_{A_m \otimes A_n} A_{m+n}^\psi \cong S \otimes_{A_n \otimes A_m} A_{m+n},$$

where A_k^ψ , $k \in \mathbb{N}$, denotes A_k , considered as an (A_k, A_k) -bimodule with the right action twisted by ψ_k , and where S is $A_n \otimes A_m$ considered as an $(A_m \otimes A_n, A_n \otimes A_m)$ -module via the obvious right multiplication and with left action given by $(a_1 \otimes a_2, s) \mapsto (a_2 \otimes a_1)s$ for $s \in S$, $a_1 \in A_m$, $a_2 \in A_n$.

For $k \in \mathbb{N}$, let 1_k denote the identity element of A_k and 1_k^ψ denote this same element considered as an element of A_k^ψ . It is straightforward to show that the map between the bimodules in (4.3) given by

$$(1_m^\psi \otimes 1_n^\psi) \otimes 1_{m+n}^\psi \mapsto (1_n \otimes 1_m) \otimes 1_{m+n}.$$

(and extended by linearity) is a well defined isomorphism. \square

Suppose A is a Hecke-like tower of algebras. If $d \neq 0$ in Definition 4.1, then $T_i(T_i - c)/d = (T_i - c)T_i/d = 1$ and so T_i is invertible for all i . It follows that T_w is invertible for all $w \in S_n$. On the other hand, if $d = 0$, then $T_i(T_i - c) = 0$ and so T_i is a zero divisor, hence not invertible. Therefore, $d \neq 0$ if and only if T_w is invertible for all w .

Lemma 4.5. *If A is a Hecke-like tower of algebras with $d \neq 0$ (equivalently, such that T_i is invertible for all i), then we have isomorphisms of functors $\nabla \cong \nabla S_{12}$ and $\Delta \cong S_{12}\Delta$ on $A\text{-mod}$ (hence also on $A\text{-pmod}$). In particular, A is dualizing.*

Proof. Let $m, n \in \mathbb{N}$ and define $w \in S_{m+n}$ by

$$w(i) = \begin{cases} m+i & \text{if } 1 \leq i \leq n, \\ i-n & \text{if } n < i \leq m+n. \end{cases}$$

Then T_w is invertible and, by Lemma 4.2(b), we have $T_w T_i = T_{w(i)} T_w$ for all $i = 1, \dots, m-1, m+1, \dots, m+n-1$. Now, let S be $A_m \otimes A_n$ considered as an $(A_n \otimes A_m, A_m \otimes A_n)$ -module via the obvious right multiplication and with left action given by $(a_1 \otimes a_2, s) \mapsto (a_2 \otimes a_1)s$ for $s \in S$, $a_1 \in A_n$, $a_2 \in A_m$. Thus, we have an isomorphism of functors $S_{12} \cong S \otimes -$. It is straightforward to verify that the map

$$A_{m+n} \rightarrow A_{m+n} \otimes_{A_n \otimes A_m} S, \quad a \mapsto a T_w \otimes (1 \otimes 1),$$

is an isomorphism of $(A_{m+n}, A_m \otimes A_n)$ -bimodules. It follows that $\nabla \cong \nabla S_{12}$. The proof that $\Delta \cong S_{12}\Delta$ is analogous. The final statement of the lemma then follows from Lemma 4.4 and Propositions 3.7 and 4.3. \square

Corollary 4.6. *All Hecke-like towers of algebras are strong and dualizing.*

Proof. It follows immediately from Lemma 4.4 and Propositions 3.7 and 4.3 that A is strong and that it is dualizing if and only if $\mathcal{K}(A)$ is cocommutative. By [BLL12, Prop. 6.1], A is isomorphic to either the tower of nilCoxeter algebras, the tower of Hecke algebras at a generic parameter, the tower of Hecke algebras at a root of unity, or the tower of 0-Hecke algebras. (While the statement of [BLL12, Prop. 6.1] is over \mathbb{C} , the proof is valid over an arbitrary algebraically closed field.) For the tower of Hecke algebras at a generic parameter or a root

of unity, we have $d \neq 0$ in Definition 4.1 and so $\mathcal{K}(A)$ is cocommutative by Lemma 4.4. For the other two towers, we will see in Sections 5 and 8 that $\mathcal{K}(A)$ is cocommutative. \square

Remark 4.7. In this section, the assumption that \mathbb{F} is algebraically closed was only used to conclude that axiom (TA4) of Definition 3.1 is satisfied and in the proof of Corollary 4.6. If \mathbb{F} is not algebraically closed but the bilinear form (3.4) is still a perfect pairing, then all the results of this section remain true except that one must replace Corollary 4.6 by the statement that the tower of nilCoxeter algebras, the tower of Hecke algebras at a generic parameter, the tower of Hecke algebras at a root of unity, and the tower of 0-Hecke algebras are all strong and dualizing.

5. NILCOXETER ALGEBRAS

In this section we specialize the constructions of Sections 2 and 3 to the tower of nilcoxeter algebras of type A . We will see that we recover Khovanov's categorification of the polynomial representation of the Weyl algebra (see [Kho01]). We let \mathbb{F} be an arbitrary field.

Definition 5.1 (Nilcoxeter algebra). The *nilcoxeter algebra* N_n is the unital \mathbb{F} -algebra generated by u_1, \dots, u_{n-1} subject to the relations

$$\begin{aligned} u_i^2 &= 0 \text{ for } i = 1, 2, \dots, n-1, \\ u_i u_j &= u_j u_i \text{ for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1, \\ u_i u_{i+1} u_i &= u_{i+1} u_i u_{i+1} \text{ for } i = 1, 2, \dots, n-2. \end{aligned}$$

The representation theory of N_n is straightforward. (We refer the reader to [Kho01] for proofs of the facts stated here.) Up to isomorphism, there is one simple module L_n , which is one dimensional, and on which the generators u_i all act by zero. The projective cover of L_n is $P_n = N_n$, considered as an N_n -module by left multiplication. We have isomorphisms of Hopf algebras

$$\begin{aligned} \mathcal{K}(N) &\cong \mathbb{Z}[x], & [P_n] &\mapsto x^n, \\ \mathcal{G}(N) &\cong \mathbb{Z}[x, x^2/2!, x^3/3!, \dots], & [L_n] &\mapsto x^n/n!. \end{aligned}$$

In both cases, the coproduct is given by $\Delta(x) = x \otimes 1 + 1 \otimes x$. We also have

$$\mathcal{G}_{\text{proj}}(N) \cong \mathbb{Z}[x],$$

and the Cartan map $\mathcal{K}(N) \rightarrow \mathcal{G}(N)$ of Definition 3.9 corresponds to the natural inclusion $\mathbb{Z}[x] \hookrightarrow \mathbb{Z}[x, x^2/2!, x^3/3!, \dots]$.

The inner product satisfies $\langle x^m, \frac{x^n}{n!} \rangle = \langle [P_m], [L_n] \rangle = \delta_{mn}$. Therefore $x^* \left(\frac{x^m}{m!} \right) = \frac{x^{m-1}}{(m-1)!}$, i.e. $x^* = \partial_x$ corresponds to partial derivation by x . Therefore the algebra \mathfrak{h} in this setting is the subalgebra of $\text{End } \mathbb{Z}[x, x^2/2!, x^3/3!, \dots]$ generated by $x, x^2!, x^3/3!, \dots$ and ∂_x . In addition, $\mathfrak{h}_{\text{proj}}$ is the algebra generated by x and ∂_x , with relation $[\partial_x, x] = 1$. The Fock space $\mathcal{F}_{\text{proj}}$ is the representation of $\mathfrak{h}_{\text{proj}}$ given by its natural action on $\mathbb{Z}[x]$. It follows from the above that $\mathbb{Q} \otimes_{\mathbb{Z}} \mathfrak{h} \cong \mathbb{Q} \otimes_{\mathbb{Z}} \mathfrak{h}_{\text{proj}}$ is the rank one Weyl algebra.

By Remark 4.7, the tower N is strong and dualizing. Thus, Theorem 3.18 provides a categorification of the polynomial representation of the Weyl algebra. In fact, (3.14) specializes to the main result of [Kho01] if one takes M and P to be the trivial A_1 -modules.

Indeed, with these choices we have $\Psi^{\otimes 2} \Delta \Psi^{-1}(P) = (\mathbb{F}_0 \otimes \mathbb{F}_1) \oplus (\mathbb{F}_1 \otimes \mathbb{F}_0)$, where \mathbb{F}_i denotes the trivial A_i -module for $i = 0, 1$. Then (3.14) becomes

$$\begin{aligned} \text{Res}_{A_n}^{A_{n+1}} \circ \text{Ind}_{A_n}^{A_{n+1}} &\cong \left(\text{Ind}_{A_{n-1}}^{A_n} \circ \text{Res}_{(\mathbb{F}_0 \otimes \mathbb{F}_1)}(\mathbb{F}_1 \otimes -) \right) \oplus \left(\text{Res}_{(\mathbb{F}_1 \otimes \mathbb{F}_0)}(M \otimes -) \right) \\ &\cong \left(\text{Ind}_{A_{n-1}}^{A_n} \circ \text{Res}_{A_{n-1}}^{A_n} \right) \oplus \text{Id}, \end{aligned}$$

which is the categorification of the relation $\partial_x x = x \partial_x + 1$ appearing in [Kho01, (13)]. (Note that while [Kho01] works over the field \mathbb{Q} , the arguments go through over more general \mathbb{F} .)

6. HECKE ALGEBRAS AT GENERIC PARAMETERS

In this section we specialize the constructions of Sections 2 and 3 to the tower of algebras corresponding to the Hecke algebras of type A at a generic parameter. The results of this section also apply to the group algebra of the symmetric group (the case when $q = 1$).

6.1. The Hecke algebra and symmetric functions. Let A_n be the Hecke algebra at a generic value of q . More precisely, assume $q \in \mathbb{C}^\times$ is not a nontrivial root of unity and let A_n be the unital \mathbb{C} -algebra with generators T_i , $i = 1, \dots, n-1$, and relations

$$\begin{aligned} T_i^2 &= q + (q-1)T_i \text{ for } i = 1, 2, \dots, n-1, \\ T_i T_j &= T_j T_i \text{ for } i, j = 1, \dots, n-1 \text{ such that } |i-j| > 1, \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1} \text{ for } i = 1, 2, \dots, n-2. \end{aligned}$$

By convention, we set $A_0 = A_1 = \mathbb{C}$. Then $A = \bigoplus_{n \in \mathbb{N}} A_n$ is a Hecke-like tower of algebras. It is well known that a complete set of irreducible A_n -modules is given by $\{S_\lambda \mid \lambda \in \mathcal{P}(n)\}$, where S_λ is the Specht module corresponding to the partition λ (see [DJ86, §6]). Since the A_n are semisimple, we have $\mathcal{K}(A) = \mathcal{G}(A)$. In fact, both are isomorphic (as Hopf algebras) to Sym , the algebra of symmetric functions in countably many variables x_1, x_2, \dots over \mathbb{Z} . This isomorphism is given by the map sending $[S_\lambda]$ to s_λ , the Schur function corresponding to the partition λ (see, for example, [Zel81]). Recall that Sym is a graded connected Hopf algebra:

$$\text{Sym} = \bigoplus_{n \geq 0} \text{Sym}_n,$$

where Sym_n is the \mathbb{Z} -submodule of Sym consisting of homogeneous polynomials of degree n . We adopt the convention that $\text{Sym}_n = 0$ for $n < 0$. The inner product (3.4) corresponds to the usual inner product on Sym under which the Schur functions are self-dual. Furthermore, the monomial and homogeneous symmetric functions are dual to each other:

$$\langle m_\lambda, h_\mu \rangle = \delta_{\lambda, \mu}, \quad \lambda, \mu \in \mathcal{P}.$$

Under this inner product, Sym is self-dual as a Hopf algebra. In other words, (Sym, Sym) is a dual pair of Hopf algebras.

6.2. The Heisenberg algebra. Applying the construction of Section 2.2 to the dual pair (Sym, Sym) , we obtain the *Heisenberg algebra* $\mathfrak{h} = \mathfrak{h}(\text{Sym}, \text{Sym})$. We obtain a minimal presentation of \mathfrak{h} by considering two collections of polynomial generators for Sym (one for Sym viewed as H^+ and one for Sym viewed as H^-). In particular, if we choose the power

sum symmetric functions p_n , $n \in \mathbb{N}_+$, in both cases, we get the usual presentation of the Heisenberg algebra:

$$[p_n, p_k] = 0, \quad [p_n^*, p_k^*] = 0, \quad [p_n^*, p_k] = n\delta_{n,k}, \quad n, k \in \mathbb{N}_+.$$

However, the p_n^*, p_n , $n \in \mathbb{N}$, are only a generating set for $\mathfrak{h} \otimes_{\mathbb{Z}} \mathbb{Q}$ since the power sum symmetric functions only generate the ring of symmetric functions over \mathbb{Q} .

On the other hand, if we choose the elementary symmetric functions e_n , $n \in \mathbb{N}_+$, and the complete symmetric functions h_n , $n \in \mathbb{N}_+$, we have the following relations:

$$(6.1) \quad [e_n, e_k] = 0, \quad [h_n^*, h_k^*] = 0, \quad [h_n^*, e_k] = e_{k-1}h_{n-1}^*, \quad n, k \in \mathbb{N}_+.$$

This gives a presentation of \mathfrak{h} (one does not need to tensor with \mathbb{Q}) and is the one used in the categorification of \mathfrak{h} given in [Kho, LS13] (for an overview, see [LS12]).

Other choices of polynomial generators result in different presentations. For the sake of completeness we record the other nontrivial relations:

$$(6.2) \quad [e_n^*, h_k] = h_{k-1}e_{n-1}^*, \quad [h_n^*, h_k] = \sum_{i \geq 1} h_{k-i}h_{n-i}^* \quad \text{and} \quad [e_n^*, e_k] = \sum_{i \geq 1} e_{k-i}e_{n-i}^*.$$

To prove these relations, we use the fact (see, for example, [Zab, Prop. 3.6]) that, for $k, n \in \mathbb{N}$, we have

$$\begin{aligned} h_k^*(h_n) &= h_{n-k}, & h_k^*(e_n) &= (\delta_{k0} + \delta_{k1})e_{n-k}, & h_k^*(p_n) &= \delta_{kn} + \delta_{k0}p_n, \\ e_k^*(h_n) &= (\delta_{k0} + \delta_{k1})h_{n-k}, & e_k^*(e_n) &= e_{n-k}, & e_k^*(p_n) &= (-1)^{k-1}\delta_{kn} + \delta_{k0}p_n, \\ p_k^*(h_n) &= h_{n-k}, & p_k^*(e_n) &= (-1)^{k-1}e_{n-k}, & p_k^*(p_n) &= n\delta_{nk} + \delta_{k0}p_n. \end{aligned}$$

Then, for example, since $\Delta(e_n) = \sum_{i=0}^n e_i \otimes e_{n-i}$, we have, by Lemma 2.8,

$$e_n^*(h_k f) = \sum_{i=0}^n e_i^*(h_k) e_{n-i}^*(f) = h_k e_n^*(f) + h_{k-1} e_{n-1}^*(f) \quad \text{for all } f \in \text{Sym}.$$

Thus $[e_n^*, h_k] = h_{k-1}e_{n-1}^*$. The other relations are proven similarly.

6.3. Categorification. By Corollary 4.6, the tower A is strong and dualizing. For $n \in \mathbb{N}$, let E_n (resp. L_n) be the one-dimensional representation of A_n on which each T_i acts by -1 (resp. by q). Then $\Delta(E_n) \cong \sum_{i=0}^n E_i \otimes E_{n-i}$ and $\Delta(L_n) \cong \sum_{i=0}^n L_i \otimes L_{n-i}$. Since $\text{Hom}_{A_n}(L_n, E_n) = 0$ unless $n = 0$ or $n = 1$ (in which case E_n and L_n are both the trivial module), we have, by (3.14),

$$\text{Res}_{L_n} \circ \text{Ind}_{E_k} \cong \nabla \left(\bigoplus_{i=0}^n \text{Res}_{L_i \otimes L_{n-i}}(E_k \otimes -) \right) \cong (\text{Ind}_{E_k} \circ \text{Res}_{L_n}) \oplus (\text{Ind}_{E_{k-1}} \circ \text{Res}_{L_{n-1}}),$$

which is a categorification of the last relation of (6.1) since, under the isomorphism $\mathcal{G}(A) \cong \mathcal{K}(A) \cong \text{Sym}$, the class of the representation L_n corresponds to h_n and the class of E_k corresponds to e_k . By (3.12), (3.13), and Lemma 4.5, we have

$$\begin{aligned} \text{Ind}_{E_n} \circ \text{Ind}_{E_k} &\cong \text{Ind}_{\nabla(E_n \otimes E_k)} \cong \text{Ind}_{\nabla(E_k \otimes E_n)} \cong \text{Ind}_{E_k} \circ \text{Ind}_{E_n}, \quad \text{and} \\ \text{Res}_{L_n} \circ \text{Res}_{L_k} &\cong \text{Res}_{\nabla(L_n \otimes L_k)} \cong \text{Res}_{\nabla(L_k \otimes L_n)} \cong \text{Res}_{L_k} \circ \text{Res}_{L_n}, \end{aligned}$$

which categorifies the first two relations of (6.1).

Remark 6.1. While we have chosen to show how Theorem 3.18 recovers a categorification of the relations (6.1), we could just have easily used it to recover categorifications of the relations (6.2). This is an illustration of the fact that Theorem 3.18 does not rely on a particular presentation of the Heisenberg double $\mathfrak{h}(A)$.

Remark 6.2. The special case of Theorem 2.11(c) for the dual pair (Sym, Sym) is known as the *Stone–von Neumann Theorem*.

Remark 6.3. Since we have $\mathcal{K}(A) = \mathcal{G}(A)$, it follows that $\mathfrak{h}_{\text{proj}} = \mathfrak{h}$ in this case (see Definition 3.13).

7. HECKE ALGEBRAS AT ROOTS OF UNITY

We now consider Hecke algebras at a root of unity. Fix $\ell \in \mathbb{N}_+$ and consider the unital \mathbb{C} -algebra A_n with generators and relations as in Section 6.1, but with q replaced by a fixed ℓ th root of unity ζ . By Corollary 4.6, $A = \bigoplus_{n \in \mathbb{N}} A_n$ is a strong dualizing tower of algebras. We refer the reader to [LLT96, §3.3] for an overview of some of the facts about Grothendieck groups stated in this section.

Let $\mathcal{J}_\ell \subseteq \text{Sym}$ be the ideal generated by the power sum symmetric functions $p_\ell, p_{2\ell}, p_{3\ell}, \dots$, and let \mathcal{J}_ℓ^\perp be its orthogonal complement relative to the standard inner product on Sym (see Section 6.1). Then there are isomorphisms of Hopf algebras

$$\mathcal{K}(A) \cong \mathcal{J}_\ell^\perp \quad \text{and} \quad \mathcal{G}(A) \cong \text{Sym}/\mathcal{J}_\ell.$$

Moreover, under these identifications, the inner product between $\mathcal{K}(A)$ and $\mathcal{G}(A)$ is that induced by the standard inner product on Sym .

Recall that a partition λ is said to be ℓ -regular if each part appears fewer than ℓ times. The specialization \bar{S}_λ of the Specht module S_λ , $\lambda \in \mathcal{P}(n)$, to $q = \zeta$ is, in general, no longer an irreducible A_n -module. However, it was shown in [DJ86, §6] that if λ is ℓ -regular, then \bar{S}_λ contains a unique maximal submodule $\text{rad } \bar{S}_\lambda$. As λ varies over the ℓ -regular partitions of n , $D_\lambda := \bar{S}_\lambda / \text{rad } \bar{S}_\lambda$ varies over a complete set of nonisomorphic irreducible representations of A_n . It follows that a basis of $\mathcal{G}(A)$ (resp. $\mathcal{K}(A)$) is given by the $[D_\lambda]$ (resp. $[P_\lambda]$, where P_λ is the projective cover of D_λ) as λ varies over the set of ℓ -regular partitions. In theory, one could compute the relations in $\mathfrak{h}(A)$ in these bases by using the results of [LLT96] to express the basis elements in terms of the standard symmetric functions and then use the relations in Section 6.2. In this way, one would obtain a presentation of $\mathfrak{h}(A)$. Of course, in general, this presentation would be far from minimal.

In fact, it turns out that $\mathfrak{h}(A)$ is an integral form of the usual Heisenberg algebra $\mathfrak{h}(\text{Sym}, \text{Sym})$ (see Section 6.2). This can be seen as follows. Recall that the set of power sum functions p_λ , $\lambda \in \mathcal{P}$, is an orthogonal basis of $\text{Sym}_\mathbb{Q} = \mathbb{Q} \otimes_{\mathbb{Z}} \text{Sym}$. (Throughout we use a subscript \mathbb{Q} to denote extension of scalars to the rational numbers.) Therefore $\mathcal{J}_{\ell, \mathbb{Q}}$ has a basis given by the set

$$\{p_\lambda \mid \ell \text{ divides } \lambda_i \text{ for at least one } i\},$$

and $\mathcal{J}_{\ell, \mathbb{Q}}^\perp$ has a basis

$$\{p_\lambda \mid \ell \text{ does not divide } \lambda_i \text{ for any } i\}.$$

Similarly, $(\text{Sym}/\mathcal{J}_\ell)_\mathbb{Q}$ has a basis

$$\{p_\lambda + \mathcal{J}_\ell \mid \ell \text{ does not divide } \lambda_i \text{ for any } i\}.$$

Remark 7.1. We see from the above that $\mathcal{G}(A_n)$ and $\mathcal{K}(A_n)$ have bases indexed, on the one hand, by the set of ℓ -regular partitions of n and, on the other hand, by the set of partitions of n in which no part is divisible by ℓ . A correspondence between these two sets of partitions is given by Glaisher’s Theorem (see, for example, [Leh46, p. 538]).

For $m \in \mathbb{N}_+$ such that ℓ does not divide m , let $q_m = p_m + \mathcal{J}_\ell$. Then we have algebra isomorphisms

$$\begin{aligned} \mathcal{J}_{\ell, \mathbb{Q}}^\perp &\cong \mathbb{Q}[p_m \mid m \in \mathbb{N}_+, \ell \text{ does not divide } m], \quad \text{and} \\ (\text{Sym}/\mathcal{J}_\ell)_{\mathbb{Q}} &\cong \mathbb{Q}[q_m \mid m \in \mathbb{N}_+, \ell \text{ does not divide } m]. \end{aligned}$$

Thus, $\mathfrak{h}(A)_{\mathbb{Q}}$ is generated by $\{p_m, q_m \mid m \in \mathbb{N}_+, \ell \text{ does not divide } m\}$ subject to the relations $[p_m, p_n] = [q_m, q_n] = 0$ and $[p_m, q_n] = m\delta_{m,n}1$. It follows that $\mathfrak{h}(A)_{\mathbb{Q}}$ is isomorphic as an algebra to the classical Heisenberg algebra $\mathfrak{h}(\text{Sym}, \text{Sym})_{\mathbb{Q}}$. Thus we have the following proposition.

Proposition 7.2. *The Heisenberg double associated to the tower of Hecke algebras at a root of unity is an integral form of the classical Heisenberg algebra:*

$$\mathfrak{h}(A)_{\mathbb{Q}} \cong \mathfrak{h}(\text{Sym}, \text{Sym})_{\mathbb{Q}}.$$

By Proposition 7.2, as ℓ varies over the positive integers, we obtain a family of integral forms of the classical Heisenberg algebra. It would be interesting to work out minimal presentations of these integral forms over \mathbb{Z} . Furthermore, the Cartan map $\mathcal{K}(A) \rightarrow \mathcal{G}(A)$ is known to have a nonzero determinant (see [BK02, Cor. 1]). Therefore, it induces an isomorphism $\mathcal{G}_{\text{proj}}(A)_{\mathbb{Q}} \cong \mathcal{G}(A)_{\mathbb{Q}}$, which implies that $\mathfrak{h}_{\text{proj}}(A)_{\mathbb{Q}} \cong \mathfrak{h}(A)_{\mathbb{Q}}$. It is not known whether $\mathfrak{h}_{\text{proj}}(A) \cong \mathfrak{h}(A)$.

It is known that the category A -pmod yields a categorification of the basic representation of $\widehat{\mathfrak{sl}}_n$ via i -induction and i -restriction functors (see [LLT96, p. 218]). Theorem 3.18 provides a categorification of the principle Heisenberg subalgebra of $\widehat{\mathfrak{sl}}_n$.

8. 0-HECKE ALGEBRAS

We now specialize the constructions of Sections 2 and 3 to the tower of 0-Hecke algebras of type A . We begin by recalling some basic facts about the rings of quasisymmetric and noncommutative symmetric functions. We refer the reader to [LMvW] for further details.

8.1. The quasisymmetric functions. Let QSym be the algebra of *quasisymmetric functions* in the variables x_1, x_2, \dots over \mathbb{Z} . Recall that this is the subalgebra of $\mathbb{Z}[[x_1, x_2, \dots]]$ consisting of shift invariant elements. That is, $f \in \text{QSym}$ if and only if, for all $k \in \mathbb{N}_+$, the coefficient in f of the monomial $x_1^{n_1} x_2^{n_2} \cdots x_k^{n_k}$ is equal to the coefficient of the monomial $x_{i_1}^{n_1} x_{i_2}^{n_2} \cdots x_{i_k}^{n_k}$ for all strictly increasing sequences of positive integers $i_1 < i_2 < \cdots < i_k$ and all $n_1, n_2, \dots, n_k \in \mathbb{N}$. The algebra QSym is a graded algebra:

$$\text{QSym} = \bigoplus_{n \geq 0} \text{QSym}_n,$$

where QSym_n is the \mathbb{Z} -submodule of QSym consisting of homogeneous elements of degree n . We adopt the convention that $\text{QSym}_n = 0$ for $n < 0$.

The algebra QSym has a basis consisting of the *monomial quasisymmetric functions* M_α , which are indexed by compositions $\alpha = (\alpha_1, \dots, \alpha_r) \in \mathcal{C}$:

$$M_\alpha = \sum_{i_1 < \dots < i_r} x_{i_1}^{\alpha_1} \cdots x_{i_r}^{\alpha_r}.$$

We adopt the convention that $M_\emptyset = 1$.

The algebra QSym has another important basis, the *fundamental quasisymmetric functions* F_α , which are defined as follows. For two compositions α, β , write $\beta \preceq \alpha$ if β is a refinement of α . For example, $(1, 2, 1) \preceq (1, 3)$. Then set

$$F_\alpha = \sum_{\beta \preceq \alpha} M_\beta, \quad \alpha \in \mathcal{C}.$$

The algebra QSym is, in fact, a graded connected Hopf algebra. To describe the coproduct, we introduce a bit of notation relating to compositions. For two compositions $\alpha = (\alpha_1, \dots, \alpha_r)$ and $\beta = (\beta_1, \dots, \beta_s)$ let $\alpha \cdot \beta = (\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_s)$ and $\alpha \odot \beta = (\alpha_1, \dots, \alpha_r + \beta_1, \dots, \beta_s)$. So, for example, if $\alpha = (1, 2, 1)$ and $\beta = (3, 5)$, then $\alpha \cdot \beta = (1, 2, 1, 3, 5)$ and $\alpha \odot \beta = (1, 2, 4, 5)$. Then the coproduct on QSym is given by either of the two following formulas:

$$\begin{aligned} \Delta(M_\alpha) &= \sum_{\alpha = \beta \cdot \gamma} M_\beta \otimes M_\gamma, \\ \Delta(F_\alpha) &= \sum_{\alpha = \beta \cdot \gamma \text{ or } \alpha = \beta \odot \gamma} F_\beta \otimes F_\gamma. \end{aligned}$$

Note that naturally $\text{Sym} \subseteq \text{QSym}$. In particular, the monomial symmetric functions can be handily expressed in terms of the monomial quasisymmetric functions:

$$(8.1) \quad m_\lambda = \sum_{\tilde{\alpha} = \lambda} M_\alpha, \quad \text{where } \tilde{\alpha} \text{ is the partition obtained by sorting } \alpha.$$

8.2. The noncommutative symmetric functions. Define NSym , the algebra of *noncommutative symmetric functions*, to be the free associative algebra (over \mathbb{Z}) generated by the alphabet $\mathbf{h}_1, \mathbf{h}_2, \dots$. Thus NSym has a basis given by $\mathbf{h}_\alpha := \mathbf{h}_{\alpha_1} \cdots \mathbf{h}_{\alpha_r}$, $\alpha \in \mathcal{C}$. This is a graded algebra:

$$\text{NSym} = \bigoplus_{n \geq 0} \text{NSym}_n,$$

where $\text{NSym}_n = \text{Span}\{\mathbf{h}_\alpha \mid \alpha \in \mathcal{C}(n)\}$. We adopt the convention that $\text{NSym}_n = 0$ for $n < 0$.

The *noncommutative ribbon Schur functions* \mathbf{r}_α are defined to be

$$\mathbf{r}_\alpha = \sum_{\alpha \preceq \beta} (-1)^{\ell(\alpha) - \ell(\beta)} \mathbf{h}_\beta, \quad \alpha \in \mathcal{C}.$$

These basis elements multiply nicely:

$$\mathbf{h}_\alpha \mathbf{h}_\beta = \mathbf{h}_{\alpha \cdot \beta} \quad \text{and} \quad \mathbf{r}_\alpha \mathbf{r}_\beta = \mathbf{r}_{\alpha \cdot \beta} + \mathbf{r}_{\alpha \odot \beta}.$$

In fact, NSym is a graded connected Hopf algebra. The coproduct is given by the formula

$$(8.2) \quad \Delta(\mathbf{h}_n) = \sum_{i=0}^n \mathbf{h}_i \otimes \mathbf{h}_{n-i}.$$

8.3. The 0-Hecke algebra and its Grothendieck groups. Let \mathbb{F} be an arbitrary field and let A_n be the unital \mathbb{F} -algebra with generators and relations as in Section 6.1, but with q replaced by 0 (i.e. the 0-Hecke algebra). Consider the tower of algebras $A = \bigoplus_{n \in \mathbb{N}} A_n$. The irreducible A_n -modules are all one-dimensional and are naturally enumerated by the set $\mathcal{C}(n)$ of compositions of n (see [Nor79, §3] and [KT97, §5.2]). Let L_α be the irreducible module corresponding to the composition $\alpha \in \mathcal{C}(n)$ and let P_α be its projective cover. We then have (see [KT97, Cor. 5.8 and Cor. 5.11] – while the statements there are for the case that $\mathbb{F} = \mathbb{C}$, the proofs remain valid over more general fields)

$$(8.3) \quad H^- = \mathcal{K}(A) \cong \text{NSym}, \quad [P_\alpha] \mapsto \mathbf{r}_\alpha,$$

$$(8.4) \quad H^+ = \mathcal{G}(A) \cong \text{QSym}, \quad [L_\alpha] \mapsto F_\alpha.$$

We also have

$$\mathcal{G}_{\text{proj}}(A) \cong \text{Sym},$$

and the Cartan map $\mathcal{K}(A) \rightarrow \mathcal{G}_{\text{proj}}(A)$ of Definition 3.9 corresponds to the projection of Hopf algebras

$$(8.5) \quad \chi: \text{NSym} \rightarrow \text{Sym}, \quad \mathbf{h}_\alpha \mapsto h_{\tilde{\alpha}}.$$

Alternatively, it is given by $\chi(\mathbf{r}_\alpha) = r_\alpha$, where r_α is the usual ribbon Schur function. This is a reformulation of [KT97, Prop. 5.9].

The bilinear form (3.4) becomes the well-known perfect Hopf pairing of the Hopf algebras QSym and NSym given as follows:

$$\begin{aligned} \langle \cdot, \cdot \rangle : \text{NSym} \times \text{QSym} &\rightarrow \mathbb{Z}, \\ \langle \mathbf{h}_\alpha, M_\beta \rangle &= \delta_{\alpha\beta} = \langle \mathbf{r}_\alpha, F_\beta \rangle, \quad \alpha, \beta \in \mathcal{C}. \end{aligned}$$

In this way, $(\text{NSym}, \text{QSym})$ is a dual pair of Hopf algebras.

8.4. The quasi-Heisenberg algebra. We now apply the construction of Section 2.2 to the dual pair $(\text{QSym}, \text{NSym})$.

Definition 8.1 ((Projective) quasi-Heisenberg algebra). We call $\mathfrak{q} := \mathfrak{h}(\text{QSym}, \text{NSym})$ the *quasi-Heisenberg algebra*. We define the *projective quasi-Heisenberg algebra* $\mathfrak{q}_{\text{proj}}$ to be the subalgebra of \mathfrak{q} generated by NSym and $\text{Sym} \subseteq \text{QSym}$ (see Definition 3.13).

Lemma 8.2. *In \mathfrak{q} we have, for all $\alpha = (\alpha_1, \dots, \alpha_r) \in \mathcal{C}$, $n \in \mathbb{N}_+$,*

$$[{}^R\mathbf{h}_n^*, M_\alpha] = M_{(\alpha_1, \dots, \alpha_{r-1})} {}^R\mathbf{h}_{n-\alpha_r}^*,$$

with the understanding that ${}^R\mathbf{h}_k^* = 0$ for $k < 0$.

Proof. By Lemma 2.8 and (8.2), we have

$${}^R\mathbf{h}_n^*(M_\alpha G) = \sum_{i=0}^n {}^R\mathbf{h}_i^*(M_\alpha) {}^R\mathbf{h}_{n-i}^*(G).$$

So, if $n \geq \alpha_r$, we have ${}^R\mathbf{h}_n^*(M_\alpha G) = M_\alpha {}^R\mathbf{h}_n^*(G) + M_{(\alpha_1, \dots, \alpha_{r-1})} {}^R\mathbf{h}_{n-\alpha_r}^*(G)$. The result follows. \square

Corollary 8.3. *For $n \in \mathbb{N}$ and $\lambda \in \mathcal{P}$, we have*

$$(8.6) \quad [{}^R\mathbf{h}_n^*, m_\lambda] = \sum_{j=1}^n m_{\lambda-j} {}^R\mathbf{h}_{n-j}^*,$$

where $\lambda - j$ is equal to the partition obtained from removing a part j from λ if λ has such a part and $m_{\lambda-j}$ is defined to be zero otherwise. In particular, for $n, k \in \mathbb{N}$, we have

$$[{}^R\mathbf{h}_n^*, p_k] = {}^R\mathbf{h}_{n-k}^*, \quad [{}^R\mathbf{h}_n^*, e_k] = e_{k-1} {}^R\mathbf{h}_{n-1}^*, \quad [{}^R\mathbf{h}_n^*, h_k] = \sum_{j=1}^n h_{k-j} {}^R\mathbf{h}_{n-j}^*.$$

Proof. Equation (8.6) follows from (8.1) and Lemma 8.2. The remainder of the relations then follow by expressing p_k , e_k and h_k in terms of the monomial symmetric functions m_λ . \square

Corollary 8.4. *The quasi-Heisenberg algebra \mathfrak{q} is generated by the set*

$$\{M_\alpha, {}^R\mathbf{h}_n^* \mid \alpha \in \mathcal{C}, n \in \mathbb{N}_+\}.$$

The M_α multiply as in QSym (for a precise description of this product, see [LMvW, §3.3.1]) and

$$[{}^R\mathbf{h}_n^*, M_\alpha] = M_{(\alpha_1, \dots, \alpha_{r-1})} {}^R\mathbf{h}_{n-\alpha_r}^*, \quad \alpha = (\alpha_1, \dots, \alpha_r) \in \mathcal{C}, \quad n \in \mathbb{N}_+.$$

Remark 8.5. We also note that the algebra \mathfrak{q} has generators

$$\{F_\alpha, {}^R\mathbf{h}_n^* \mid \alpha \in \mathcal{C}, n \in \mathbb{N}_+\}.$$

From a representation theoretic point of view, these are more natural since the F_α correspond to simple A_n -modules (see (8.4)). The F_α then multiply as in QSym (for a precise description of this product, see [LMvW, §3.3.1]) and

$$(8.7) \quad [{}^R\mathbf{h}_n^*, F_\alpha] = \sum_{i=1}^{\alpha_r} F_{(\alpha_1, \dots, \alpha_{r-i})} {}^R\mathbf{h}_{n-i}^*, \quad \alpha = (\alpha_1, \dots, \alpha_r) \in \mathcal{C}, n \in \mathbb{N}_+.$$

Remark 8.6. Note that the above presentations are far from minimal. There are polynomial generators of QSym, given by modified Lyndon words (see [DS99, Th. 1.5]), which one could use instead of the M_α in the above presentation. This would result in a minimal presentation of \mathfrak{q} .

The following result gives a presentation of the projective quasi-Heisenberg algebra in terms of generators and relations.

Proposition 8.7. *The algebra $\mathfrak{q}_{\text{proj}}$ is generated by the set*

$$\{e_n, {}^R\mathbf{h}_n^* \mid n \in \mathbb{N}\}.$$

The relations are

$$[e_n, e_k] = 0, \quad [{}^R\mathbf{h}_n^*, e_k] = e_{k-1} {}^R\mathbf{h}_{n-1}^*, \quad n, k \in \mathbb{N}.$$

Proof. This follows immediately from the definition of $\mathfrak{q}_{\text{proj}}$ and Corollary 8.3. \square

Remark 8.8. Note the similarity of the presentation of Proposition 8.7 to the presentation of the usual Heisenberg algebra $\mathfrak{h}(\text{Sym}, \text{Sym})$ given in (6.1). The only difference is that the h_n^* commute, whereas the ${}^R\mathbf{h}_n^*$ do not. There is a natural surjective map of algebras $\mathfrak{q}_{\text{proj}} \rightarrow \mathfrak{h}(\text{Sym}, \text{Sym})$ given by $e_n \mapsto e_n, {}^R\mathbf{h}_n^* \mapsto h_n^*, n \in \mathbb{N}_+$.

8.5. Fock spaces and categorification. As described in Section 2.3, the quasi-Heisenberg algebra \mathfrak{q} acts naturally on QSym and we call this the *lowest weight Fock space representation* of \mathfrak{q} . By Theorem 2.11(c), any representation of \mathfrak{q} generated by a lowest weight vacuum vector is isomorphic to QSym.

Similarly, as in Definition 3.14, the projective quasi-Heisenberg algebra $\mathfrak{q}_{\text{proj}}$ acts naturally on Sym and we call this the *lowest weight Fock space representation* of $\mathfrak{q}_{\text{proj}}$. As a $\mathfrak{q}_{\text{proj}}$ -module, Sym is generated by the lowest weight vacuum vector $1 \in \text{Sym}$. By Proposition 3.15(c), any representation of $\mathfrak{q}_{\text{proj}}$ generated by a lowest weight vacuum vector is isomorphic to Sym. However, this representation is not faithful since it factors through the projection from $\mathfrak{q}_{\text{proj}}$ to the usual Heisenberg algebra (see Remark 8.8). On the other hand, the highest weight Fock space representation of $\mathfrak{q}_{\text{proj}}$ is faithful (see Proposition 3.16).

By Remark 4.7, A is a strong dualizing tower of algebras. Therefore, Theorem 3.18 yields a categorification of the Fock space representations of \mathfrak{q} and $\mathfrak{q}_{\text{proj}}$. For instance, it is straightforward to verify that

$$\begin{aligned} \Delta(L_\alpha) &\cong \bigoplus_{\alpha=\beta \cdot \gamma \text{ or } \alpha=\beta \odot \gamma} L_\beta \otimes L_\gamma \quad \text{for all } \alpha \in \mathcal{C}, \\ \Psi^{\otimes 2} \Delta \Psi^{-1}(P_{(n)}) &\cong \Delta(P_{(n)}) \cong \bigoplus_{i=0}^n P_{(i)} \otimes P_{(n-i)} \quad \text{for all } n \in \mathbb{N}. \end{aligned}$$

For $\alpha = (\alpha_1, \dots, \alpha_r) \in \mathcal{C}$ and $i \in \{0, \dots, \alpha_r\}$, it follows that $\text{Res}_{P(i)} L_\alpha = L_{(\alpha_1, \dots, \alpha_r - i)}$. Thus we have

$$\text{Res}_{P(n)} \circ \text{Ind}_{L_\alpha} \cong \nabla \left(\bigoplus_{i=0}^n \text{Res}_{P(i) \otimes P(n-i)} (L_\alpha \otimes -) \right) \cong \bigoplus_{i=0}^{\alpha_r} \text{Ind}_{L_{(\alpha_1, \dots, \alpha_r - i)}} \text{Res}_{P(n-i)},$$

which is a categorification of the relation (8.7). The categorification of the multiplication of the elements F_α , $\alpha \in \mathcal{C}$, follows from the computation of the induction in $A\text{-mod}$ (see, for example, the proof of [DKKT97, Prop. 4.15]).

9. APPLICATION: QSym IS FREE OVER Sym

As a final application of the methods of the current paper, we use the generalized Stone–von Neumann Theorem for $\mathfrak{q}_{\text{proj}}$ (Proposition 3.15) to prove that QSym is free over Sym. This gives a proof that is quite different from the traditional one using modified Lyndon words (see [DS99, Cor. 1.6]).

Lemma 9.1. *Suppose V is a $\mathfrak{q}_{\text{proj}}$ -module which is generated (as a $\mathfrak{q}_{\text{proj}}$ -module) by a finite set of lowest weight vacuum vectors. Then V is a direct sum of copies of lowest weight Fock space.*

Proof. Let $\{v_i\}_{i \in I}$ denote a set of lowest weight vacuum vectors that generates V and such that I has minimal cardinality. We claim that

$$(9.1) \quad \mathbb{Z}v_i \cap \mathbb{Z}v_j = \{0\} \quad \text{for all } i \neq j.$$

Suppose, on the contrary, that $\mathbb{Z}v_i \cap \mathbb{Z}v_j \neq \{0\}$ for some $i \neq j$. Then $n_i v_i = n_j v_j$ for some $n_i, n_j \in \mathbb{Z}$. Let $m = \gcd(n_i, n_j)$ and choose $a_i, a_j \in \mathbb{Z}$ such that $m = a_i n_i + a_j n_j$. Set $w = a_j v_i + a_i v_j$. Then w is clearly a lowest weight vacuum vector, and we have

$$\frac{n_i}{m} w = \frac{1}{m} (a_j n_i v_i + a_i n_i v_j) = \frac{1}{m} (a_j n_j v_j + a_i n_i v_j) = v_j.$$

Similarly, $\frac{n_j}{m} w = v_i$. Thus $\{v_k\}_{k \in I \setminus \{i, j\}} \cup \{w\}$ is a set of lowest weight vacuum vectors that generates V , contradicting the minimality of the cardinality of I .

By Proposition 3.15(c), $\mathfrak{q}_{\text{proj}} \cdot v_i \cong \text{Sym}$ as $\mathfrak{q}_{\text{proj}}$ -modules. It then follows from Proposition 3.15(a) and (9.1) that $\mathfrak{q}_{\text{proj}} \cdot v_i \cap \mathfrak{q}_{\text{proj}} \cdot v_j = \{0\}$ for $i \neq j$. The lemma follows. \square

Define an increasing filtration of $\mathfrak{q}_{\text{proj}}$ -submodules of QSym as follows. For $n \in \mathbb{N}$, let

$$\text{QSym}^{(n)} := \sum_{\ell(\alpha) \leq n} \mathfrak{q}_{\text{proj}} \cdot M_\alpha.$$

In particular, note that $\text{QSym}^{(0)} = \text{Sym}$. We adopt the convention that $\text{QSym}^{(-1)} = \{0\}$.

Proposition 9.2. *The space QSym of quasisymmetric functions is free as a Sym-module.*

Proof. Note that, for $\alpha \in \mathcal{C}$ such that $\ell(\alpha) = n$, we have ${}^R \mathbf{h}_m^*(M_\alpha) \in \text{QSym}^{(n-1)}$ for any $m > 0$. Therefore, in the quotient $V_n = \text{QSym}^{(n)} / \text{QSym}^{(n-1)}$, such M_α are lowest weight vacuum vectors. It is clear that these vectors generate V_n , and therefore, by Lemma 9.1,

$$V_n = \bigoplus_{v \in \mathcal{S}_n} \text{Sym} \cdot v,$$

where \mathcal{S}_n is some collection of vacuum vectors in V_n .

Consider the short exact sequence

$$0 \rightarrow \text{QSym}^{(n-1)} \rightarrow \text{QSym}^{(n)} \rightarrow V_n \rightarrow 0.$$

Since V_n is a free (hence projective) Sym-module, the above sequence splits. Therefore, if $\text{QSym}^{(n-1)}$ is free over Sym, then so is $\text{QSym}^{(n)}$.

By the argument in the previous paragraph we can choose nested sets of vectors in QSym

$$\tilde{\mathcal{S}}_0 \subseteq \tilde{\mathcal{S}}_1 \subseteq \tilde{\mathcal{S}}_2 \subseteq \cdots$$

such that, for every $n \in \mathbb{N}$, we have $\text{QSym}^{(n)} = \bigoplus_{\tilde{v} \in \tilde{\mathcal{S}}_n} \text{Sym} \cdot \tilde{v}$. Let $\tilde{\mathcal{S}} = \bigcup_{n \in \mathbb{N}} \tilde{\mathcal{S}}_n$. Then

$$\text{QSym} = \bigoplus_{v \in \tilde{\mathcal{S}}} \text{Sym} \cdot v. \quad \square$$

REFERENCES

- [ARS95] Maurice Auslander, Idun Reiten, and Sverre O. Smalø. *Representation theory of Artin algebras*, volume 36 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995.
- [BK02] Jonathan Brundan and Alexander Kleshchev. Cartan determinants and Shapovalov forms. *Math. Ann.*, 324(3):431–449, 2002.
- [BL09] Nantel Bergeron and Huilan Li. Algebraic structures on Grothendieck groups of a tower of algebras. *J. Algebra*, 321(8):2068–2084, 2009.
- [BLL12] Nantel Bergeron, Thomas Lam, and Huilan Li. Combinatorial Hopf algebras and towers of algebras—dimension, quantization and functorality. *Algebr. Represent. Theory*, 15(4):675–696, 2012.
- [CL12] Sabin Cautis and Anthony Licata. Heisenberg categorification and Hilbert schemes. *Duke Math. J.*, 161(13):2469–2547, 2012.
- [CR81] Charles W. Curtis and Irving Reiner. *Methods of representation theory. Vol. I*. John Wiley & Sons Inc., New York, 1981.
- [DJ86] Richard Dipper and Gordon James. Representations of Hecke algebras of general linear groups. *Proc. London Math. Soc. (3)*, 52(1):20–52, 1986.
- [DKKT97] Gérard Duchamp, Alexander Klyachko, Daniel Krob, and Jean-Yves Thibon. Noncommutative symmetric functions. III. Deformations of Cauchy and convolution algebras. *Discrete Math. Theor. Comput. Sci.*, 1(1):159–216, 1997. Lie computations (Marseille, 1994).
- [DS99] E. J. Ditters and A. C. J. Scholtens. Free polynomial generators for the Hopf algebra QSym of quasisymmetric functions. *J. Pure Appl. Algebra*, 144(3):213–227, 1999.
- [FJW00] Igor B. Frenkel, Naihuan Jing, and Weiqiang Wang. Vertex representations via finite groups and the McKay correspondence. *Internat. Math. Res. Notices*, (4):195–222, 2000.
- [Gei77] Ladnor Geissinger. Hopf algebras of symmetric functions and class functions. In *Combinatoire et représentation du groupe symétrique (Actes Table Ronde C.N.R.S., Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976)*, pages 168–181. Lecture Notes in Math., Vol. 579. Springer, Berlin, 1977.
- [Haz08] M. Hazewinkel, editor. *Handbook of algebra. Vol. 5*, volume 5 of *Handbook of Algebra*. Elsevier/North-Holland, Amsterdam, 2008.
- [Kho] Mikhail Khovanov. Heisenberg algebra and a graphical calculus. arXiv:1009.3295.
- [Kho01] Mikhail Khovanov. Nilcoxeter algebras categorify the Weyl algebra. *Comm. Algebra*, 29(11):5033–5052, 2001.
- [KT97] Daniel Krob and Jean-Yves Thibon. Noncommutative symmetric functions. IV. Quantum linear groups and Hecke algebras at $q = 0$. *J. Algebraic Combin.*, 6(4):339–376, 1997.
- [Leh46] D. H. Lehmer. Two nonexistence theorems on partitions. *Bull. Amer. Math. Soc.*, 52:538–544, 1946.
- [LLT96] Alain Lascoux, Bernard Leclerc, and Jean-Yves Thibon. Hecke algebras at roots of unity and crystal bases of quantum affine algebras. *Comm. Math. Phys.*, 181(1):205–263, 1996.
- [LMvW] K. Luoto, S. Mykytiuk, and S. van Willigenburg. An introduction to quasisymmetric Schur functions. Available at <http://www.math.ubc.ca/~steph/papers/QuasiSchurBook.pdf>.

- [LS12] Anthony Licata and Alistair Savage. A survey of Heisenberg categorification via graphical calculus. *Bull. Inst. Math. Acad. Sin. (N.S.)*, 7(2):291–321, 2012.
- [LS13] Anthony Licata and Alistair Savage. Hecke algebras, finite general linear groups, and Heisenberg categorification. *Quantum Topol.*, 4(2):125–185, 2013.
- [Lu94] Jiang-Hua Lu. On the Drinfel’d double and the Heisenberg double of a Hopf algebra. *Duke Math. J.*, 74(3):763–776, 1994.
- [Nor79] P. N. Norton. 0-Hecke algebras. *J. Austral. Math. Soc. Ser. A*, 27(3):337–357, 1979.
- [STS94] M. A. Semenov-Tian-Shansky. Poisson Lie groups, quantum duality principle, and the quantum double. In *Mathematical aspects of conformal and topological field theories and quantum groups (South Hadley, MA, 1992)*, volume 175 of *Contemp. Math.*, pages 219–248. Amer. Math. Soc., Providence, RI, 1994.
- [SY11] Andrzej Skowroński and Kunio Yamagata. *Frobenius algebras. I*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2011. Basic representation theory.
- [Zab] Mike Zabrocki. Introduction to symmetric functions. Available at <http://garsia.math.yorku.ca/ghana03/chapters/mainfile2.pdf>.
- [Zel81] Andrey V. Zelevinsky. *Representations of finite classical groups*, volume 869 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1981.

A. SAVAGE: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA
URL: <http://mysite.science.uottawa.ca/asavag2/>
E-mail address: alistair.savage@uottawa.ca

O. YACOBI: SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY
URL: <http://www.maths.usyd.edu.au/u/oyacobi/>
E-mail address: oyacobi@maths.usyd.edu.au